COPYRIGHT© 2019 EDIZIONI MINERVA MEDICA

This document is protected by international copyright laws. No additional reproduction is authorized. It is permitted for personal use to download and save only one file and print only one copy of this Article. It is not permitted to make additional copies (either sporadically or systematically, either printed or electronic) of the Article for any purpose. It is not permitted to distribute the electronic copy of the article through online internet and/or intranet file sharing systems, electronic mailing or any other means which may allow access to the Article. The use of all or any part of the Article for any Commercial Use is not permitted. The creation of derivative works from the Article is not permitted. The production of reprints for personal or commercial use is not permitted. It is not permitted to remove, cover, overlay, obscure, block, or change any copyright notices or terms of use which the Publisher may post on the Article. It is not permitted to frame or use framing techniques to enclose any trademark, logo, or other proprietary information of the Publisher.

© 2018 EDIZIONI MINERVA MEDICA
Online version at http://www.minervamedica.it

European Journal of Physical and Rehabilitation Medicine 2019 April;55(2):231-40
DOI: 10.23736/S1973-9087.18.04782-2

ORIGINAL ARTICLE

Brace treatment of Idiopathic Scoliosis is effective for a curve over 40 degrees, but is the evaluation of Cobb angle the only parameter for the indication of treatment?

Angelo G. AULISA [1] *, Vincenzo GUZZANTI [1, 2], Francesco FALCIGLIA [1], Marco GIORDANO [1], Marco GALLI [3], Lorenzo AULISA [3]

[1]Unit of Orthopedics and Traumatology, Bambino Gesù Children's Hospital, Institute of Scientific Research, Rome, Italy; [2]University of Cassino, Cassino, Frosinone, Italy; [3]Department of Orthopedics, "Agostino Gemelli" University Hospital, Catholic University of the Sacred Heart School of Medicine, Rome, Italy

*Corresponding author: Angelo G. Aulisa, Unit of Orthopedics and Traumatology, Bambino Gesù Children's Hospital, Institute of Scientific Research, P.zza S. Onofrio 4 - 00165 Rome, Italy. E-mail: angelogabriele.aulisa@fastwebnet.it

ABSTRACT

BACKGROUND: The recent literature showed positive results for bracing of patients with idiopathic scoliosis above 45° who refused surgery. However, no one has investigated whether other parameters can affect the results.

AIM: The aim of this study was to evaluate the effectiveness of bracing in idiopathic scoliosis with curves above 40° and to verify the mechanical and biological parameters which go beyond the simple bend value expressed in Cobb degrees.

DESIGN: This is an observational controlled cohort study nested in a prospective clinical on-going database including 1,238 patients with idiopathic scoliosis.

SETTING: Inpatients and outpatients in Rome.

POPULATION: The study enrolled 160 patients with idiopathic scoliosis with curves above 40°.

METHODS: This is a prospective study based on an ongoing database including 1,238 patients with idiopathic scoliosis. The patients studied had idiopathic scoliosis with curves of 40° or more, Risser grade 0-4, and had refused any surgical treatment. 160 patients met the inclusion criteria. Of these, 104 patients had a definite outcome, 28 abandoned treatment and 28 are currently under treatment. The minimum duration of follow-up was 24 months. X-rays were used to obtain Cobb degrees and torsion of the apical vertebrae (Perdriolle's method). Three outcomes were distinguished according to SRS-SOSORT criteria: correction, stabilization and progression. To achieve the second aim, we divided the sample into subgroups according to Cobb degrees (<45°; ≥45°), Risser (0-2; 3-4) and rotation (<20; ≥20). Furthermore, logistic regression was applied by Stepwise Regression.

RESULTS: The results of our study showed that in 104 patients with a definite outcome the Cobb mean value was initially 47±5.25 SD and 34.19 ±8.45 SD at follow-up. Perdriolle was initially 20.04±5.53 SD and 16.76±7.04 at follow-up. Overall, 81 patients (78%) obtained a curve correction, and stabilization was achieved in 14 cases (13%). Nine patients experienced curve progression (9%), 16 patients were recommended for surgery because the curve at follow up was over 45°. The analysis of subgroups shows that with Cobb <45° at baseline, the average reduction was 11.46° Cobb, while in cases with Cobb ≥45 at baseline, the mean correction was 13.74° Cobb. In subgroups with Perdriolle <20° at baseline, the average reduction was 16.02° Cobb, while in cases with Perdriolle ≥20° at baseline, the mean correction was 8.4° Cobb. In subgroups with Risser 0-2 at baseline, the average reduction was 14.7° Cobb, while in cases with Risser 3-4 at baseline, the mean correction was 6.7° Cobb. The logistic regression model shows significance for the initial value of Perdriolle and Risser.

CONCLUSIONS: Our results indicate that an adequate conservative treatment must definitely be considered for patients with scoliotic curves who refuse surgery; the results will be better particularly if the rotation is lower than 20 and Risser is between 0-2.

CLINICAL REHABILITATION IMPACT: With the simultaneous evaluation of the Cobb angle, the vertebral rotation and the potential vertebral growth, it was possible to predict the final results at the start of treatment.

(*Cite this article as:* Aulisa AG, Guzzanti V, Falciglia F, Giordano M, Galli M, Aulisa L. Brace treatment of Idiopathic Scoliosis is effective for a curve over 40 degrees, but is the evaluation of Cobb angle the only parameter for the indication of treatment? Eur J Phys Rehabil Med 2019;55:231-40. DOI: 10.23736/S1973-9087.18.04782-2)

KEY WORDS: Scoliosis; Brace; Conservative treatment.

The specific pathological features of adolescent idiopathic scoliosis (AIS) are characterized by a process of evolution of the initial conditions that, if not treated, determines serious consequences such as curve progression, lower back pain, psychosocial problems and minor cardiopulmonary consequences.[1-6]

COPYRIGHT© 2019 EDIZIONI MINERVA MEDICA

This document is protected by international copyright laws. No additional reproduction is authorized. It is permitted for personal use to download and save only one file and print only one copy of this Article. It is not permitted to make additional copies (either sporadically or systematically, either printed or electronic) of the Article for any purpose. It is not permitted to distribute the electronic copy of the article through online internet and/or intranet file sharing systems, electronic mailing or any other means which may allow access to the Article. The use of all or any part of the Article for any Commercial Use is not permitted. The creation of derivative works from the Article is not permitted. The production of reprints for personal or commercial use is not permitted. It is not permitted to remove, cover, overlay, obscure, block, or change any copyright notices or terms of use which the Publisher may post on the Article. It is not permitted to frame or use framing techniques to enclose any trademark, logo, or other proprietary information of the Publisher.

The aim of the conservative treatment in AIS, usually performed in growing patients, is to prevent the progression of the curve.

In the older literature several authors propose that conservative treatment does not represent a truly effective option but only a procrastination of surgery, aimed at limiting the worsening of the curves while waiting for the right age for surgical treatment. Goldberg *et al.* underline that in the North American population, the incidence of surgery in patients treated by standard bracing is the same as that in observation-only patients.[7-9]

Despite the lack of data supporting brace treatment, the positivists stated, "The truth is that an effective method of early non-surgical treatment is available. That method is bracing."[10] Despite this, several studies attesting to the effectiveness of the brace have been published in recent years by members of the International Society on Scoliosis Orthopaedic and Rehabilitation Treatment (SOSORT).[11-17] Weinstein in the BrAIST Study showed that, in the North American Population, the rate of treatment success was 72% after bracing, compared with 48% after observation only.[18]

The reasons for the controversy can, still today, be summarized as follows:

• our insufficient knowledge about the etiopathogenetic factors of scoliosis hinders the search for a causal therapy for idiopathic scoliosis;

• although the scoliotic curves have a potentially worsening evolution, this is a probabilistic event and the result is due from both biological and mechanical factors;

• the results of conservative treatment are not always the same in the sense that, in relation to the methods and the therapeutic regime employed, the treatment is not completed in all cases. Finally, the possibility that orthopedic treatments can achieve a recovery of the curve, albeit a partial one, is contested;

• there is still insufficient knowledge about the biomechanical factors that affect both the worsening evolution process and the treatment of the curves. This has led to the empiricism that often characterizes the indication for the type of brace, its manufacture and the clinical management of the different types of treatment.[19]

However, the indication to conservative treatment for scoliosis with more than 40° Cobb degree curves remains controversial, even if there have been recent reports of some positive results in bracing patients with idiopathic scoliosis above 45° who refused surgical treatment.[20-23]

The aims of the study were:

• evaluate the effectiveness of bracing in idiopathic scoliosis with curves over 40° Cobb, following the guidelines on standards of management of idiopathic scoliosis (SOSORT);[24]

• verify the mechanical and biological parameters which go beyond the simple bend value expressed in Cobb degrees, and which could be used as indicators to conservative treatment even in cases that are usually included in the "gray area".

Materials and methods

Design

This is an observational controlled cohort study nested in a prospective clinical on-going database including 1238 patients with idiopathic scoliosis. Patients were recruited both in public hospitals and in private clinics. Inclusion criteria were: idiopathic scoliosis, at least one curve of 40° or more, Risser stage 0-4, age over 10 years old, and had firmly refused any surgical treatment.

Furthermore, the study was conducted using the STROBE guidelines (STROBE Statement—Checklist of items that should be included in reports of cohort studies).

Population

The study was carried out during May 2014. From a database including 1238 patients, we found 160 patients meeting the inclusion criteria; of these, 104 patients had a definite outcome, 28 abandoned treatment and 28 are currently under treatment. The minimum duration of follow-up was 24 months after the end of treatment.

All patients gave written consent for their clinical data management to be used for research purposes; the study respects the statements of the requirements of the Helsinki Declaration concerning medical experimentation on human beings.

Bracing

Patients with thoracolumbar and lumbar curves were prescribed the Progressive Action Short Brace (PASB), while the Lyon brace was prescribed for those with thoracic or double curves. Full-time bracing was prescribed for all patients (*i.e.*, max 22 hours daily, min 20 hours daily). The daily number of hours of bracing was defined for each patient according to clinical needs and acceptance. In order to maximize compliance to treatment, patients were always followed by the same doctor. Furthermore, controls were performed every 2 months. Frequent checks allowed verification and implementation of compliance, creating an open and friendly doctor-patient relationship. Close checks were

COPYRIGHT© 2019 EDIZIONI MINERVA MEDICA

This document is protected by international copyright laws. No additional reproduction is authorized. It is permitted for personal use to download and save only one file and print only one copy of this Article. It is not permitted to make additional copies (either sporadically or systematically, either printed or electronic) of the Article for any purpose. It is not permitted to distribute the electronic copy of the article through online internet and/or intranet file sharing systems, electronic mailing or any other means which may allow access to the Article. The use of all or any part of the Article for any Commercial Use is not permitted. The creation of derivative works from the Article is not permitted. The production of reprints for personal or commercial use is not permitted. It is not permitted to remove, cover, overlay, obscure, block, or change any copyright notices or terms of use which the Publisher may post on the Article. It is not permitted to frame or use framing techniques to enclose any trademark, logo, or other proprietary information of the Publisher.

also performed to maximize bracing effectiveness over time. Weaning was started when the beginning of ring-apophysis fusion could be seen on a lateral (LL) radiograph view,[25] corresponding to a Risser sign 4 or 5 on an anterior-posterior (AP) standing radiograph view. Weaning consisted of 2 to 4 hours bracing reduction at 2-month intervals.

Endpoints

For the present study, only X-rays performed at conventional times were considered: start of treatment (t1), 4-6 months after start of treatment (t2), intermediate time between t1 and t4 (t3), end of treatment (t4), 2-year minimum follow-up from end of treatment (t5).

For each patient, AP and LL view standing X-rays of the whole spine were performed. X-rays before treatment (t1) as well as those at t4 and t5 were taken without brace. All other radiographic controls were performed with the patient wearing the brace, in order to assess the corrective action of bracing. The first X-ray was obtained at 4-6 months from the beginning of treatment. All other controls were performed once a year. All radiographs were taken at our Institute, at a distance of 2 meters, using a 36×91 cm film. The AP view was used to determine the patient's skeletal age (the European Risser's sign) and to obtain the curve magnitude (CM: Cobb's method) and torsion of the apical vertebra (TA: Perdriolle's method).[26, 27] Two independent observers carried out the measurements. The end vertebrae were preselected by the senior author (LA) to reduce inter-observer error, but the end-vertebrae can change from one X-ray to another. Curves were classified according to Scoliosis Research Society (SRS) into thoracic, thoracolumbar, lumbar, and double major. SRS-SOSORT criteria for bracing studies were applied.[28] As recommended by the SRS Committee on Bracing and Non-operative Management, outcomes were classified as follows: 1) correction (percentage of patients with ≤5° curve progression); 2) stabilization (final Cobb angle not differing more than 5º from its initial value); 3) progression (percentage of patients with ≥5° progression at maturity); and 4) percentage of patients with curves exceeding 45° at maturity and of those recommended for or who had undergone surgery. To achieve the second aim we have also divided the sample into subgroups according to Cobb degrees (<45°; ≥45°), Risser (0-2; 3-4) and rotation (<20; ≥20).

Statistical analysis

Statistical analysis was performed using the Graphpad Prism 6.0.

For all variables, normality of data was verified by the Kolmogorov-Smirnov Test. Results were analyzed in relation to CM t5-t1 at follow-up, assuming that CM t5-t1 was not within the Cobb's method±5 range of error.[27] Changes in CM and TA over time from t1 through t5 were assessed via one-way analysis of variance (ANOVA) for repeated measures. Tukey's post-test was applied when needed. The model was fitted for age, type of curve, and type of bracing. All analyses were performed according to the intention-to-treat principle. Missing data at follow-up were managed according to the Last Observation Carried Forward (LOCF) method. All tests were two-sided, with significance set at $P<0.05$. Results are presented as mean±standard deviation (SD).

Logistic regression

Logistic regression was used to determine predictively the individual probability of success in each case using the Statistix 9 program.

A reduction at the last control of more than 11 Cobb degrees compared to the initial value was considered a success (variable =1); a smaller reduction, stabilization or worsening was considered a failure (variable =0).

Based on this subdivision, the 104 cases studied showed: successes 54 cases; failures 50 cases, prevalence of successes 53.8%.

Logistic regression was applied by Stepwise Regression, assuming as dependent variable success (0 or 1) and as independent starting variables: sex, age, duration of treatment, initial Cobb and Perdriolle degrees, Risser Index.

Results


Analyses of patients with a definite outcome

A definite outcome was recorded for 104 patients, 96 females (93%) and 8 males (7%), mean age 12.88±1.86 years and 17.92±1.65 years at t1 and t4, respectively. The mean duration of treatment was 63.12±17.44 months, with an average length of follow-up of 60.86±57.85 (range 24-276) months.

Curve type distribution was as follows: thoracic (N.=43; 41.3%), thoracolumbar (N.=33; 31.7%), lumbar (N.=12; 11%), double (N.=11; 10.5%) and double thoracic (N.=5; 4.8%). The Progressive Action Short Brace (PASB) was prescribed in 57 patients, and the Lyon brace in 47 patients.


Changes in CM over time were statistically significant (P for trend <0.0001) (Figure 1), with a mean val-

COPYRIGHT© 2019 EDIZIONI MINERVA MEDICA



Tukey's multiple comparison test	Mean diff	Q	Adjusted P value	95% CI of diff
t1 *vs.* t2	16.23	20.57	<0.0001	13.18 to 19.29
t1 *vs.* t3	19.44	24.64	<0.0001	16.39 to 22.50
t1 *vs.* t4	18.46	23.23	<0.0001	15.39 to 21.54
t1 *vs.* t5	12.83	15.89	<0.0001	9.703 to 15.96
t2 *vs.* t3	3.212	4.069	0.0338	0.1561 to 6.267
t2 *vs.* t4	2.234	2.810	0.2738	-0.8441 to 5.312
t2 *vs.* t5	-3.401	4.211	0.0252	-6.528 to -0.2740
t3 *vs.* t4	-0.9775	1.230	0.9080	-4.056 to 2.101
t3 *vs.* t5	-6.613	8.187	<0.0001	-9.740 to -3.486
t4 *vs.* t5	-5.635	6.928	<0.0001	-8.784 to -2.486

Figure 1.—Changes in curve magnitude in Cobb degrees from the beginning of treatment (t1) to 2-year minimum follow-up from end of weaning (t5). Each box depicts the interquartile range, with the median indicated by the black center line. Error bars show the data distribution, with the whiskers corresponding to the minimum and maximum values.



Tukey's multiple comparisons test	Mean diff	Q	Adjusted P value	95% CI of diff
t1 *vs.* t2	2.767	4.373	0.0178	0.3169 to 5.216
t1 *vs.* t3	3.907	6.176	0.0001	1.458 to 6.357
t1 *vs.* t4	4.910	7.642	< 0.0001	2.422 to 7.397
t1 *vs.* t5	3.278	5.089	0.0032	0.7839 to 5.772
t2 *vs.* t3	1.141	1.799	0.7087	-1.315 to 3.596
t2 *vs.* t4	2.143	3.328	0.1302	-0.3503 to 4.636
t2 *vs.* t5	0.5114	0.7921	0.9807	-1.989 to 3.011
t3 *vs.* t4	1.002	1.556	0.8063	-1.491 to 3.496
t3 *vs.* t5	-0.6293	0.9747	0.9588	-3.129 to 1.871
t4 *vs.* t5	-1.632	2.490	0.3979	-4.169 to 0.9055

Figure 2.—Changes in apical torsion in Perdriolle degrees from the beginning of treatment (t1) to 2-year minimum follow-up from end of weaning (t5). Each box depicts the interquartile range, with the median indicated by the black center line. Error bars show the data distribution, with the whiskers corresponding to the minimum and maximum values.

ue of 47.02±5.25° Cobb at start of treatment (t1) and 34.19±8.45° Cobb at follow-up (t5). A less significant pattern was observed for TA (P for trend <0.01) (Figure 2), with a mean value of 20.04±5.53° Perdriolle at beginning (t1) and 16.76±7.04° Perdriolle at follow-up (t5).

Overall, 81 patients (78%) obtained a curve correction, while stabilization was achieved in 14 cases (13%). Nine patients experienced curve progression (9%), 16 patients were subsequently recommended for surgery because at follow up the curve was over 45° (Table I).

The analysis of the total sample shows a significant correlation between rotation at baseline and mean curve correction in Cobb degrees, but not for Cobb at baseline and mean curve correction in Cobb degrees (Figure 3). The

TABLE I.—*Outcomes of the subgroup.*

	Curve correction (N. %)	Curve stabilization (N. %)	Curve progression (N. %)	Surgery referrals (N. %)
<45° Cobb (N.=47)	36 (77%)	7 (14%)	4 (9%)	5 (10%)
≥45° Cobb (N.=57)	45 (78%)	5 (13%)	7 (9%)	11 (19%)
<20° Perdriolle (N.=55)	53 (96%)	1 (2%)	1 (2%)	1 (2%)
≥ 20° Perdriolle (N.=49)	28 (57%)	13 (27%)	8 (16%)	15 (30%)
≥25° Perdriolle (N.=22)	3 (14%)	15 (68%)	4 (18%)	18 (81)
Risser 0-2 (N.=68)	59 (86%)	6 (9%)	3 (5%)	7 (10%)
Risser 3-4 (N.=36)	22 (61%)	12 (33%)	2 (6%)	9 (25%)
PASB (N.=57)	47 (82%)	5 (9%)	5 (9%)	5 (9%)
Lyon Brace (N.=47)	34 (72%)	9 (19%)	4 (9%)	11 (23%)

This document is protected by international copyright laws. No additional reproduction is authorized. It is permitted for personal use to download and save only one file and print only one copy of this Article. It is not permitted to make additional copies (either sporadically or systematically, either printed or electronic) of the Article for any purpose. It is not permitted to distribute the electronic copy of the article through online internet and/or intranet file sharing systems, electronic mailing or any other means which may allow access to the Article. The use of all or any part of the Article for any Commercial Use is not permitted. The creation of derivative works from the Article is not permitted. The production of reprints for personal or commercial use is not permitted. It is not permitted to remove, cover, overlay, obscure, block, or change any copyright notices or terms of use which the Publisher may post on the Article. It is not permitted to frame or use framing techniques to enclose any trademark, logo, or other proprietary information of the Publisher.

COPYRIGHT© 2019 EDIZIONI MINERVA MEDICA




Figure 3.—Correlations between mean curve correction in Cobb degrees and Perdriolle at baseline (A) and Cobb at baseline (B).

analysis of subgroups shows that with CM<45° at t1, the average reduction was 11.46° Cobb, while in cases with CM≥45 at t1, the mean correction was 13.74° Cobb. The multiple comparisons in subgroup <45° were significant between the age at baseline and the mean curve correction in Cobb and Perdriolle degrees (Table II, III). Similar results were reported for CM≥45. Significant correlations were observed between Perdriolle at baseline and mean curve correction in Cobb degrees (Figure 4, 5).

In particular, 36 patients in subgroups with CM<45 obtained a curve correction with a mean value of 13.75°, stabilization was achieved in 7 cases (mean 1.7°) and 4 patients experienced curve progression (mean 7°). In comparison, 45 patients in subgroups with CM≥45 obtained a curve correction with a mean value of 14.88°, stabilization was achieved in 5 cases (mean 0.2°) and 7 patients experienced curve progression (mean 5.14°).

In subgroups with TA<20° at t1, the average reduction was 16.02° Cobb, while in cases with TA≥20 at t1, the mean correction was 8.4° Cobb. The multiple compari-

TABLE II.—*Comparisons between different variables in Cobb subgroups (C_M<45).*

Tukey's multiple comparisons test	Mean diff	*t*	Adjusted P value	95% CI of diff
Cobb t1 *vs.* Cobb t5	11.46	3.950	0.0137	0.7542 to 22.16
CM t5-t1 *vs.* TA t5-t1	-8.099	2.711	0.4437	-19.12 to 2.923
Age at baseline *vs.* CM t5-t1	23.37	8.103	< 0.0001	12.73 to 34.02
Risser *vs.* CM t5-t1	12.50	4.334	< 0.0001	1.858 to 23.15
P t1 *vs.* P t5	2.287	0.7928	0.9982	-8.357 to 12.93
Age at t1 *vs.* TA t5-t1	15.28	5.140	< 0.0001	4.308 to 26.24
Risser *vs.* TA t5-t1	4.403	1.481	0.9789	-6.564 to 15.37

TABLE III.—*Comparisons between different variables in Cobb subgroups (C_M≥45).*

Tukey's multiple comparisons test	Mean diff	*t*	Adjusted P value	95% CI of diff
Cobb t1 *vs.* Cobb t5	13.74	4.796	0.0004	3.174 to 24.30
CM t5-t1 *vs.* TA t5-t1	-10.31	3.414	0.0837	-21.45 to 0.8281
Age at t1 *vs.* CM t5-t1	26.38	9.210	<0.0001	15.82 to 36.95
Risser *vs.* CM t5-t1	15.40	5.376	<0.0001	4.838 to 25.97
P t1 *vs.* P t5	4.073	1.422	0.9982	-6.492 to 14.64
Age at t1 *vs.* TA t5-t1	20.86	7.119	<0.0001	10.06 to 31.67
Risser *vs.* TA t5-t1	5.091	1.737	0.9789	-5.716 to 15.90




Figure 4.—Correlations between mean curve correction in Cobb degrees and Risser at baseline (A) Cobb at baseline (B) and Perdriolle at baseline (C) in CM<45.

This document is protected by international copyright laws. No additional reproduction is authorized. It is permitted for personal use to download and save only one file and print only one copy of this Article. It is not permitted to make additional copies (either sporadically or systematically, either printed or electronic) of the Article for any purpose. It is not permitted to distribute the electronic copy of the article through online internet and/or intranet file sharing systems, electronic mailing or any other means which may allow access to the Article. The use of all or any part of the Article for any Commercial Use is not permitted. The creation of derivative works from the Article is not permitted. The production of reprints for personal or commercial use is not permitted. It is not permitted to remove, cover, overlay, obscure, block, or change any copyright notices or terms of use which the Publisher may post on the Article. It is not permitted to frame or use framing techniques to enclose any trademark, logo, or other proprietary information of the Publisher.

COPYRIGHT© 2019 EDIZIONI MINERVA MEDICA




Figure 5.—Correlations between mean curve correction in Cobb degrees and Risser at baseline (A) Cobb at baseline (B) and Perdriolle at baseline (C) in CM≥45.

sons in subgroups TA<20° and TA≥20° were significant between the age at baseline and the mean curve correction in Cobb and Perdriolle degrees (Table IV, V). Significant correlations were found between Perdriolle at baseline and mean curve correction in Cobb degrees (Figure 6, 7).

In subgroups with Risser 0-2 at t1, the average reduc-

TABLE IV.—*Comparisons between different variables in Perdriolle subgroups (T_A<20).*

Tukey's multiple comparisons test	Mean diff	*t*	Adjusted P value	95% CI of diff
Cobb t1 *vs.* Cobb t5	16.02	5.630	<0.0001	5.526 to 26.52
CM t5-t1 *vs.* TA t5-t1	-10.78	3.714	0.0317	-21.49 to -0.07732
Age at t1 *vs.* CM t5-t1	27.96	9.871	<0.0001	17.52 to 38.41
Risser *vs.* CM t5-t1	16.89	5.963	<0.0001	6.444 to 27.34
Perdriolle t1 *vs.* Perdriolle t5	4.094	1.439	0.9979	-6.401 to 14.59
Age at t1 *vs.* TA t5-t1	17.18	5.919	<0.0001	6.477 to 27.89
Risser *vs.* TA t5-t1	6.109	2.105	0.8713	-4.595 to 16.81

TABLE V.—*Comparisons between different variables in Perdriolle subgroups (T_A≥20).*

Tukey's multiple comparisons test	Mean diff	*t*	Adjusted P value	95% CI of diff
Cobb t1 *vs.* Cobb t5	8.400	2.964	0.2706	-2.057 to 18.86
CM t5-t1 *vs.* TA t5-t1	-7.202	2.372	0.7059	-18.41 to 4.000
Age at t1 *vs.* CM t5-t1	20.88	7.367	<0.0001	10.42 to 31.33
Risser *vs.* CM t5-t1	10.10	3.565	0.0525	-0.3543 to 20.56
Perdriolle t1 *vs.* Perdriolle t5	1.558	0.5533	>0.9999	-8.831 to 11.95
Age at t1 *vs.* TA t5-t1	13.68	4.690	0.0006	2.916 to 24.44
Risser *vs.* TA t5-t1	2.900	0.9945	>0.9999	-7.860 to 13.66




Figure 6.—Correlations between mean curve correction in Cobb degrees and Risser at baseline (A) Cobb at baseline (B) and Perdriolle at baseline (C) in TA<20.

This document is protected by international copyright laws. No additional reproduction is authorized. It is permitted for personal use to download and save only one file and print only one copy of this Article. It is not permitted to make additional copies (either sporadically or systematically, either printed or electronic) of the Article for any purpose. It is not permitted to distribute the electronic copy of the article through online internet and/or intranet file sharing systems, electronic mailing or any other means which may allow access to the Article. The use of all or any part of the Article for any Commercial Use is not permitted. The creation of derivative works from the Article is not permitted. The production of reprints for personal or commercial use is not permitted. It is not permitted to remove, cover, overlay, obscure, block, or change any copyright notices or terms of use which the Publisher may post on the Article. It is not permitted to frame or use framing techniques to enclose any trademark, logo, or other proprietary information of the Publisher.

COPYRIGHT© 2019 EDIZIONI MINERVA MEDICA




Figure 7.—Correlations between mean curve correction in Cobb degrees and Risser at baseline (A) Cobb at baseline (B) and Perdriolle at baseline (C) in TA≥20.

TABLE VI.—*Outcomes of the patients at follow-up.*

	Beginning of treatment	Follow-up
>45° Cobb	57	22
<40° Cobb	0	65
<30° Cobb	0	38

TABLE VII.—*Outcomes of the patients in agreement of SRS-SOSORT criteria.*

Outcome	N.	%
Progressed 5° or more	3	3
Unchanged	20	19.2
Improved 5° or more	81	77.8
Total	104	

TABLE VIII.—*Decision-making matrix.*

	Success	Failure
T+	50	4
T-	6	44
TOT	56	48

TABLE IX.—*Diagnostic table.*

Sensibility	89.3%
Specificity:	91.7%
Predicting value (VP) +	92.6%
Predicting value (VP) -	88%
Accuracy	90.4%




Figure 8.—ROC curves.
ROC1: Risser+Perdriolle; ROC2: Perdriolle; ROC3: Risser.

tion was 14.7° Cobb, while in cases with Risser 3-4 at t1, the mean correction was 6.7° Cobb.

Treatment outcomes are shown in Table I, VI, VII.

Logistic regression

The model discarded, for the non-significance of the coefficients, all the variables except for the initial value of Perdriolle and Risser.

It provided a regression equation for logit L: L= 10.41-0.403 * Initial Pedriolle -1.21 * Risser

The results are detailed by L: Odds= Exp(L); Probab. % = 100* Odds /(1+Odds)

Taking into account a discriminating threshold of 50% probability, the decision-making matrix can be built (Table VIII) and gives the diagnostic table (Table IX). The method is highly predictive with an accuracy of 90.4%. It appears particularly suitable for predicting positivity (*VP* +: 92.6%).

ROC analysis

To see how the result could vary according to the decision-making threshold adopted, a ROC curve was constructed using the method described by Galli *et al.*[29]

For completeness, the ROC curves were also drawn using Perdriolle and Risser as parameters.

The graph in Figure 8 shows how the curve determined with Perdriolle + Risser shows a significantly increased area under the curve (AUC= 0.961) (P<0.05) compared

This document is protected by international copyright laws. No additional reproduction is authorized. It is permitted for personal use to download and save only one file and print only one copy of this Article. It is not permitted to make additional copies (either sporadically or systematically, either printed or electronic) of the Article for any purpose. It is not permitted to distribute the electronic copy of the article through online internet and/or intranet file sharing systems, electronic mailing or any other means which may allow access to the Article. The use of all or any part of the Article for any Commercial Use is not permitted. The creation of derivative works from the Article is not permitted. The production of reprints for personal or commercial use is not permitted. It is not permitted to remove, cover, overlay, obscure, block, or change any copyright notices or terms of use which the Publisher may post on the Article. It is not permitted to frame or use framing techniques to enclose any trademark, logo, or other proprietary information of the Publisher.

COPYRIGHT© 2019 EDIZIONI MINERVA MEDICA

TABLE X.—*Shows the sensitivity values (TPF) for some threshold values (FPF= 1-Specificity) and the complementary data obtained from the two-factor curve ROC analysis.*

FPF	Sensitivity	FPF	Sensitivity	AUC	96.1%
2.5	70.3	30	97.5	ES	2%
5	82.9	35	98	Q	90.6%
10	91.1	40	98.4	DOR	92.2±1.97
15	94.2	45	98.7		
20	95.8	50	98.9		
25	96.8				

to that obtained with the individual parameters, and that they practically do not differ among themselves. Furthermore, Table X shows the sensitivity values (TPF) for some threshold values (FPF= 1-Specificity) and the complementary data obtained from the two-factor curve ROC analysis; in particular the Q point is 90.6% and the Diagnostic Odds Ratio (DOR) is over 90.

Discussion

This study confirmed that brace treatment could be useful for adolescent scoliosis with curves over 40° Cobb. The indication to surgical treatment is greatly decreased by bracing, while the probability of success (reduction of the curve of at least 5°) is deeply influenced by the severity of torsional deformity (Perdriolle degree) and skeletal maturity (Risser degree).

The efficacy of brace treatment has been questioned for many years. Recently, many European studies, in particular the Braist study that included a randomized arm, have shown the effectiveness of bracing in adolescent patients.[13, 14, 16, 23, 30] However, no adolescents with curves above 40° have been included in these studies because the only generally accepted therapeutic option for this specific population is surgery. Furthermore, in recent years societies such as SRS and SOSORT have accepted bracing as a therapeutic option in adolescents with curves of 45 degrees and the Braist Study considered progression to 50 degrees as conservative treatment failure.

The patients in our study had surgical indication but, despite our explaining to them the advantages of surgery and the poor evidence of the efficacy of conservative treatment for this type of curve, they firmly refused treatment and consequently received the conservative treatment. Nevertheless, our patients have the strongest motivation to avoid a spinal fusion, resulting in high compliance rates. Compliance is an important factor in obtaining good results with an efficacious brace.

Moreover, the results of this study are in accord with those reported by Lusini for patients with a curve over 45 degrees that showed conservative treatment failure in only 23.5% of patients,[20] However, our results provide new information about the parameters to be evaluated before starting treatments in the "gray area".

To better explain the results, the importance of the degree of torsion and the rationale of this study, we must clarify some principles of biomechanics.

In particular, studies on the brace-spine interaction and analysis of the instability of the scoliotic spine,[31] have underlined that stopping the evolutionary process of the idiopathic scoliosis and, when possible, correcting the deformity and deformations, can be achieved provided that the brace is projected according to precise mechanical requirements and that the biological structure can react in a suitable way to the induced actions.[32-34]

Studies on the biological reaction to mechanical actions could provide tools for defining the indications to and limits of conservative treatment, which could be used as well as evaluating the curve in Cobb degrees.

Previous papers observed that a deformed spine has a mobility which reduces the natural "dynamic stability".[32, 35] In a growing subject, this phenomenon induces a progressive increase of the vertebral asymmetry causing a worsening of the curve originally provoked by biological factors.[33, 36]

The mechanical behavior of the scoliotic spine clarifies the theoretical suppositions concerning brace treatment.[37]

The correction can be achieved *via* forces which can modify the model of distribution of the solicitations. Intervertebral discs are viscoelastic structures that can distort themselves and return to their normal shape. Correct position and good viscoelastic equilibrium are the basic conditions for physiological biomechanics of the intervertebral disc. In fact, a condition of hysteresis of the disc would make the disc unable to transmit the corrective actions of the brace. The twisting component is the element which characterizes the scoliotic curve.

In a scoliotic spine, vertebrae are locked in a permanent twist, which distorts the vertebra and puts the disc in a position of permanent rotation. This "patho"-mechanical

This document is protected by international copyright laws. No additional reproduction is authorized. It is permitted for personal use to download and save only one file and print only one copy of this Article. It is not permitted to make additional copies (either sporadically or systematically, either printed or electronic) of the Article for any purpose. It is not permitted to distribute the electronic copy of the article through online internet and/or intranet file sharing systems, electronic mailing or any other means which may allow access to the Article. The use of all or any part of the Article for any Commercial Use is not permitted. The creation of derivative works from the Article is not permitted. The production of reprints for personal or commercial use is not permitted. It is not permitted to remove, cover, overlay, obscure, block, or change any copyright notices or terms of use which the Publisher may post on the Article. It is not permitted to frame or use framing techniques to enclose any trademark, logo, or other proprietary information of the Publisher.

COPYRIGHT© 2019 EDIZIONI MINERVA MEDICA

This document is protected by international copyright laws. No additional reproduction is authorized. It is permitted for personal use to download and save only one file and print only one copy of this Article. It is not permitted to make additional copies (either sporadically or systematically, either printed or electronic) of the Article for any purpose. It is not permitted to distribute the electronic copy of the article through online internet and/or intranet file sharing systems, electronic mailing or any other means which may allow access to the Article. The use of all or any part of the Article for any Commercial Use is not permitted. The creation of derivative works from the Article is not permitted. The production of reprints for personal or commercial use is not permitted. It is not permitted to remove, cover, overlay, obscure, block, or change any copyright notices or terms of use which the Publisher may post on the Article. It is not permitted to frame or use framing techniques to enclose any trademark, logo, or other proprietary information of the Publisher.

setting induces an increase in the stiffness of the spine towards twisting, subsequently leading to hysteresis of the disc. This condition is related to the entity of rotation and the initial value of the G modulus, that changes in relation to its position and to the patient's age.[31]

These results are confirmed by the biomechanical analysis; in fact, over time, the elastic characteristics of the spine undergo considerable variations.

The different degrees of intervertebral disc elasticity explain the different degrees of permeability to external actions and, therefore, to the different evolutionary patterns seen in scoliosis.

In children, the lower value of the torsion modules of the discs can indeed promote a spontaneous evolution toward serious deformity. However, this evolution can be stopped by administering a proper conservative treatment and achieving an effective recovery of the deformity.

Therefore recovery, independent of the initial Cobb degrees, is subordinate to the concomitant presence of two essential conditions.

The growth of vertebrae: a limited residual growth precludes any remodelling of the vertebral geometry. If treatment is started at the end of growth, the curve correction is obtained mainly by deforming the disco-elastic structures, and in fact the correction will vanish when the patient removes the brace.

The disc alteration is located in the elastic-plastic deformity area, therefore allowing low degrees of spinal rotation. The elastic return to zero degrees, after removal of the torsional moment, can only happen if the rotation is sufficiently below the field of linear elasticity and the applied torsional moment rapidly loses its effect. Otherwise a portion of the energy furnished remain in the disc and provoke an increase in rigidity, thus producing the phenomenon of hysteresis.

This explains why, for a scoliosis with a medium grade of rotation, recovery can only be limited due to the partial deformation of the discs. For a scoliosis with a high grade of rotation, the alteration of the discs allows no correction with the brace treatment, only a stabilization of the curve.

Finally, we would like to underline the importance of the increase in the physiological values of G with the advance of age. In fact a scoliotic curve with an important rotation considered not recoverable in the adolescent may be susceptible to meaningful recovery in infants.

Limitations of the study

Our study has certain limitations: 1) some of our cases lack long-term follow-up which could show a progression requiring fusion; 2) the actual gold standard for treatment for curves over 45 degrees is fusion which excludes being able to perform a randomized controlled study and thus reduces the scientific strength of the study.

Conclusions

The indications to conservative treatment in idiopathic scoliosis are normally defined only taking into consideration the lateral deviations on the coronal plane. In fact, Cobb degrees are used to determine the "area" of therapeutic orientation with the determination of a "gray area" and a "surgical area". This type of evaluation has been confirmed as being useful for all patients where conservative treatment can be guaranteed to stop the curve progression. However, this study demonstrates the limits of an evaluation criterion that does not adequately take into consideration elements such as residual vertebral growth and the elastic potential of the discs. Their capacity to induce, during conservative treatment, a recovery of the deformity, even partial, determines the different results in the different "areas."

Although the results do not allow a precise definition of new "areas" based upon the convergence of different factors which could change the evolution of the curves, it is likely that the Risser sign and the degree of torsion presented at the first observation are primary elements when considering the indication to conservative treatment and predicting a successful outcome of the treatment. Thus, the conservative treatment of scoliosis, administered in accordance with SOSORT criteria, plays a primary role in the management of this condition, even when patients are in the "gray" or surgical areas.

Finally, this new approach could offer a less controversial interpretation of the different results obtained with similar initial Cobb degrees in different subjects receiving conservative treatment. With the simultaneous evaluation of the Cobb angle, vertebral rotation and potential vertebral growth it is possible to predict the end results at the start of treatment.

References

1. Weinstein SL, Dolan LA, Cheng JC, Danielsson A, Morcuende JA. Adolescent idiopathic scoliosis. Lancet 2008;371:1527–37.

2. Branthwaite MA. Cardiorespiratory consequences of unfused idiopathic scoliosis. Br J Dis Chest 1986;80:360–9.

3. Ascani E, Bartolozzi P, Logroscino CA, Marchetti PG, Ponte A, Savini R, *et al.* Natural history of untreated idiopathic scoliosis after skeletal maturity. Spine 1986;11:784–9.

COPYRIGHT© 2019 EDIZIONI MINERVA MEDICA

4. Pehrsson K, Bake B, Larsson S, Nachemson A. Lung function in adult idiopathic scoliosis: a 20 year follow up. Thorax 1991;46:474–8.

5. Weinstein SL, Ponseti IV. Curve progression in idiopathic scoliosis. J Bone Joint Surg Am 1983;65:447–55.

6. Padua R, Ceccarelli E, Pitra L, De Santis V, Aulisa L. Radioaerosol lung scintigraphy in idiopathic scoliosis. Stud Health Technol Inform 2002;88:279–84.

7. Negrini S, Aulisa L, Ferraro C, Fraschini P, Masiero S, Simonazzi P, *et al.* Italian guidelines on rehabilitation treatment of adolescents with scoliosis or other spinal deformities. Eura Medicophys 2005;41:183–201.

8. Weiss HR, Negrini S, Rigo M, Kotwicki T, Hawes MC, Grivas TB, *et al*. Indications for conservative management of scoliosis (SOSORT guidelines). Stud Health Technol Inform 2008;135:164–70.

9. Richards BS, Bernstein RM, D'Amato CR, Thompson GH. Standardization of criteria for adolescent idiopathic scoliosis brace studies: SRS Committee on Bracing and Nonoperative Management. Spine 2005;30:2068–75.

10. Bassett GS, Bunnell WP, MacEwen GD. Treatment of idiopathic scoliosis with the Wilmington brace. Results in patients with a twenty to thirty-nine-degree curve. J Bone Joint Surg Am 1986;68:602–5.

11. Asher MA, Burton DC. Adolescent idiopathic scoliosis: natural history and long term treatment effects. Scoliosis 2006;1:2.

12. Weiss HR, Goodall D. The treatment of adolescent idiopathic scoliosis (AIS) according to present evidence. A systematic review. Eur J Phys Rehabil Med 2008;44:177–93.

13. Aulisa AG, Guzzanti V, Galli M, Perisano C, Falciglia F, Aulisa L. Treatment of thoraco-lumbar curves in adolescent females affected by idiopathic scoliosis with a progressive action short brace (PASB): assessment of results according to the SRS committee on bracing and nonoperative management standardization criteria. Scoliosis 2009;4:21.

14. Negrini S, Atanasio S, Fusco C, Zaina F. Effectiveness of complete conservative treatment for adolescent idiopathic scoliosis (bracing and exercises) based on SOSORT management criteria: results according to the SRS criteria for bracing studies - SOSORT Award 2009 Winner. Scoliosis 2009;4:19.

15. Lonstein JE, Carlson JM. The prediction of curve progression in untreated idiopathic scoliosis during growth. J Bone Joint Surg Am 1984;66:1061–71.

16. Aulisa AG, Guzzanti V, Falciglia F, Giordano M, Marzetti E, Aulisa L. Lyon bracing in adolescent females with thoracic idiopathic scoliosis: a prospective study based on SRS and SOSORT criteria. BMC Musculoskelet Disord 2015;16:316.

17. Aulisa AG, Guzzanti V, Marzetti E, Giordano M, Falciglia F, Aulisa L. Brace treatment in juvenile idiopathic scoliosis: a prospective study in accordance with the SRS criteria for bracing studies - SOSORT award 2013 winner. Scoliosis 2014;9:3.

18. Weinstein SL, Dolan LA, Wright JG, Dobbs MB. Design of the Bracing in Adolescent Idiopathic Scoliosis Trial (BrAIST). Spine 2013;38:1832–41.

19. Little DG, Song KM, Katz D, Herring JA. Relationship of peak height velocity to other maturity indicators in idiopathic scoliosis in girls. J Bone Joint Surg Am 2000;82:685–93.

20. Lusini M, Donzelli S, Minnella S, Zaina F, Negrini S. Brace treatment is effective in idiopathic scoliosis over 45°: an observational prospective cohort controlled study. Spine J 2014;14:1951–6.

21. Negrini S, Negrini F, Fusco C, Zaina F. Idiopathic scoliosis patients with curves more than 45 Cobb degrees refusing surgery can be effectively treated through bracing with curve improvements. Spine J 2011;11:369–80.

22. Aulisa AG, Guzzanti V, Giordano M, Falciglia F, Fuiano M, Aulisa L. Conservative treatment in adolescent idiopathic scoliosis with curves over 45°: is the measurement in Cobb degrees the only parameter to be considered? Scoliosis 2014;9:1–2.

23. De Mauroy J. Prospective study of 136 adolescent scoliosis of more than 40° treated with the Lyon brace. Scoliosis 2013;8:O34.

24. Negrini S, Aulisa AG, Aulisa L, Circo AB, de Mauroy JC, Durmala J, *et al.* 2011 SOSORT guidelines: orthopaedic and Rehabilitation treatment of idiopathic scoliosis during growth. Scoliosis 2012;7:3.

25. Bick EM, Copel JW. The ring apophysis of the human vertebra; contribution to human osteogeny. II. J Bone Joint Surg Am 1951;33-A:783–7.

26. Omeroğlu H, Ozekin O, Biçimoğlu A. Measurement of vertebral rotation in idiopathic scoliosis using the Perdriolle torsionmeter: a clinical study on intraobserver and interobserver error. Eur Spine J 1996;5:167–71.

27. Morrissy RT, Goldsmith GS, Hall EC, Kehl D, Cowie GH. Measurement of the Cobb angle on radiographs of patients who have scoliosis. Evaluation of intrinsic error. J Bone Joint Surg Am 1990;72:320–7.

28. Negrini S, Hresko TM, O'Brien JP, Price N; SOSORT Boards; SRS Non-Operative Committee. Recommendations for research studies on treatment of idiopathic scoliosis: Consensus 2014 between SOSORT and SRS non-operative management committee. Scoliosis 2015;10:8.

29. Galli M, Ciriello V, Menghi A, Aulisa AG, Rabini A, Marzetti E. Joint line tenderness and McMurray tests for the detection of meniscal lesions: what is their real diagnostic value? Arch Phys Med Rehabil 2013;94:1126–31.

30. Dolan LA, Wright JG, Weinstein SL. Effects of bracing in adolescents with idiopathic scoliosis. N Engl J Med 2014;370:681.

31. Fineschi G, Aulisa L, Vinciguerra A. La rigidezza del rachide alla torsione. Progressi Patol Vertebr 1990;11:109–17.

32. Aulisa L, Vinciguerra A, Tamburrelli F, Lupparelli S, Di Legge V. Biomechanical analysis of the elastic behaviour of the spine with aging. Stud Health Technol Inform 1997;37:229–32.

33. Stokes IA, Burwell RG, Dangerfield PH; IBSE. Biomechanical spinal growth modulation and progressive adolescent scoliosis—a test of the 'vicious cycle' pathogenetic hypothesis: summary of an electronic focus group debate of the IBSE. Scoliosis 2006;1:16.

34. Clin J, Aubin CE, Parent S, Sangole A, Labelle H. Comparison of the biomechanical 3D efficiency of different brace designs for the treatment of scoliosis using a finite element model. Eur Spine J 2010;19:1169–78.

35. Aulisa L, Lupparelli S, Pola E, Aulisa AG, Mastantuoni G, Pitta L. Biomechanics of the conservative treatment in idiopathic scoliotic curves in surgical "grey-area". Stud Health Technol Inform 2002;91:412–8.

36. Stokes IA. Analysis and simulation of progressive adolescent scoliosis by biomechanical growth modulation. Eur Spine J 2007;16:1621–8.

37. Aulisa L, Di Benedetto A, Vinciguerra A. [Biomechanical analysis of the spinal brace system in idiopathic scoliosis]. Arch Putti Chir Organi Mov 1981;31:185–94.

Conflicts of interest.—The authors certify that there is no conflict of interest with any financial organization regarding the material discussed in the manuscript.

Authors' contributions.—Angelo G. Aulisa participated in the conception, design and coordination, acquisition of data, analysis and interpretation of data and drafted the manuscript; Angelo G. Aulisa and Marco Giordano performed the statistical analysis. Marco Giordano, Francesco Falciglia and Vincenzo Guzzanti helped to draft the manuscript; Lorenzo Aulisa participated in the conception, design and coordination and helped to draft the manuscript. All authors read and approved the final manuscript.

Article first published online: March 7, 2018. - Manuscript accepted: March 6, 2018. - Manuscript revised: March 1, 2018. - Manuscript received: April 11, 2017.

This document is protected by international copyright laws. No additional reproduction is authorized. It is permitted for personal use to download and save only one file and print only one copy of this Article. It is not permitted to make additional copies (either sporadically or systematically, either printed or electronic) of the Article for any purpose. It is not permitted to distribute the electronic copy of the article through online internet and/or intranet file sharing systems, electronic mailing or any other means which may allow access to the Article. The use of all or any part of the Article for any Commercial Use is not permitted. The creation of derivative works from the Article is not permitted. The production of reprints for personal or commercial use is not permitted. It is not permitted to remove, cover, overlay, obscure, block, or change any copyright notices or terms of use which the Publisher may post on the Article. It is not permitted to frame or use framing techniques to enclose any trademark, logo, or other proprietary information of the Publisher.