https://doi.org/10.3301/ABSGI.2022.02

Torino 19-21 September 2022



a cura della Società Geologica Italiana



GEOSCIENCES FOR A SUSTAINABLE FUTURE













PRESIDENTS OF THE CONGRESS

Rodolfo Carosi and Daniele Castelli.

SCIENTIFIC COMMITTEE

Germana Barone, Fabrizio Berra, Matteo Berti, Fernando Camara, Piergulio Cappelletti, Claudio Chiarabba, Rita Chirico, Renato Colucci, Chiara D'Ambrogi, Rosanna De Rosa, Laura De Santis, Daniela Ducci, Patrizia Fumagalli, Biagio Giaccio, Fausto Guzzetti, Giulia Innamorati, Ilaria Mazzini, Mimmo Palano, Claudia Piromallo, Mario Soldati, Stefania Venturi, Andrea Zanchi.

ORGANISING COMMITTEE

Donato Belmonte, Sabrina Bonetto, Bernardo Carmina, Francesco Dela Pierre, Evdokia Tema, Lorenza Fascio, Salvatore Iaccarino, Andrea Maffeis, Nadia Malaspina, Chiara Montomoli, Marcello Natalicchio, Alessandro Petroccia, Fabio Massimo Petti, Fabrizio Piana, Franco Rolfo, Licia Santoro, Mario Tribaudino, Alessandro Zuccari.

ABSTRACT BOOK EDITORS

Bernardo Carmina, Lorenza Fascio, Giulia Innamorati, Marco Pasero & Fabio Massimo Petti.

COVER IMAGE:

Aerial cityscape image of Turin during sunset.

Papers, data, figures, maps and any other material published are covered by the copyright own by the **Società Geologica** Italiana.

DISCLAIMER: The Società Geologica Italiana, the Editors are not responsible for the ideas, opinions, and contents of the papers published; the authors of each paper are responsible for the ideas opinions and contents published.

La Società Geologica Italiana, i curatori scientifici non sono responsabili delle opinioni espresse e delle affermazioni pubblicate negli articoli: l'autore/i è/sono il/i solo/i responsabile/i.

Numerical analysis of the post-seismic effects on groundwater flow after the Amatrice-Visso-Norcia 2016 seismic sequence

Zullo E.*1, Albano M.2, Saroli M.1-2, Moro M.2, Testa G.1, Bonora N.1, Petitta M.3 & Doglioni C.2-3

¹ Dipartimento di Ingegneria Civile e Meccanica, Università di Cassino e del Lazio meridionale. ² INGV, Roma. ³ Dipartimento di Scienze della Terra, Sapienza Università di Roma.

Corresponding author e-mail: enrica.zullo@unicas.it

Keywords: groundwater flow, Norcia earthquake, numerical modelling.

The hydrogeological setting of the Sibillini Mts. is mainly characterized by the coexistence of a shallow and a basal regional aquifer (Viaroli et al., 2021) whose spatial distribution, recharge area and groundwater dynamics are strongly controlled by a complex regional tectonic framework.

The 2016 seismic sequence that struck Central Italy altered the hydrodynamic response of the carbonate fractured aquifer, causing sustained variations in the groundwater flow. In particular, after the Mw 6.5, 30 October main shock (called Norcia earthquake), caused by the dislocation of the Vettore Mt. normal fault, a surplus of groundwater discharge was observed at the springs bordering the western side of the Sibillini Mts. and in the Nera River basin. Furthermore, the Torbidone spring in the Norcia plain, exhausted since 1979, reappeared after the seismic event. The observed post-seismic groundwater surplus was attributed to a variation of the hydraulic gradient and an increase in bulk permeability probably caused by the fracture cleaning effect, which caused a shift of the groundwater divide after the fault rupture (Mastrorillo et al., 2020). Such phenomenon induced an eastward shift of the piezometric divide so that the springs located on the eastern side of the Sibillini Mts. strongly reduced their flow, thus causing several problems in the groundwater supply system.

In this work, we aim at assessing the effect of the coseismic dislocation of the Vettore fault on the crustal stress and strain field, the consequent modification of the hydraulic properties of the shallow and basal regional aquifers of the Sibillini Mts. and the related groundwater flow variations. For this purpose, we collected geological and hydrogeological data from the available literature and used them to analyse the area affected by the 30 October earthquake with 2D and 3D numerical models. We first created a simplified 2D geometric model at surface scale (up to 2 km deep), including the spatial relationship between the main tectonic elements and the boundaries of the main hydrogeological complexes, to model the groundwater flow before and after the seismic event. Then a regional-scale 3D model is developed, whose hydraulic parameters are calibrated as a function of the stress and strain changes caused by Vettore fault rupture to obtain two different hydrological scenarios: pre- and post-faulting.

Mastrorillo L., Saroli M., Viaroli S., Banzato F., Valigi D. & Petitta M. (2020) - Sustained post-seismic effects on groundwater flow in fractured carbonate aquifers in Central Italy. Hydrological Processes, 34(5), 1167-1181.

Viaroli S., Mirabella F., Mastrorillo L., Angelini S. & Valigi D. (2021) - Fractured carbonate aquifers of Sibillini Mts. (Central Italy). Journal of Maps, 17(2), 140-149.

