

Review article

Inflammation as a mediator between adverse childhood experiences and adult depression: A meta-analytic structural equation model

Andrea Zagaria^a, Valeria Fiori^a, Mariacarla Vacca^a, Caterina Lombardo^a, Carmine M. Pariante^b, Andrea Ballesio^{a,*}

^a Department of Psychology, Faculty of Medicine and Psychology, Sapienza University of Rome, Italy

^b Department of Psychological Medicine, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, UK

ARTICLE INFO

Keywords:

Trauma
Early life stress
Biomarker
Immune system
Meta-analysis
Depression
BMI
MASEM

ABSTRACT

Exposure to adverse childhood experiences (ACEs) confers a higher risk of developing depression in adulthood, yet the mediation of inflammation remains under debate. To test this model, we conducted a systematic review and two-stage structural equation modelling meta-analysis of studies reporting correlations between ACEs before age 18, inflammatory markers and depression severity in adulthood. Scopus, Pubmed, Medline, PsycInfo, and CINAHL were searched up to 2 October 2023. Twenty-two studies reporting data on C-reactive protein (CRP, $n = 12,935$), interleukin-6 (IL-6, $n = 4108$), tumour necrosis factor- α (TNF- α , $n = 2256$) and composite measures of inflammation ($n = 1674$) were included. Unadjusted models revealed that CRP ($\beta = 0.003$, 95 % LBCI 0.0002 to 0.0068), IL-6 ($\beta = 0.003$, 95 % LBCI 0.001 to 0.006), and composite inflammation ($\beta = 0.009$, 95 % LBCI 0.004 to 0.018) significantly mediated the association between ACEs and adult depression. The mediation effects no longer survived after adjusting for BMI; however, a serial mediation model revealed that BMI and IL-6 sequentially mediated the association between ACEs and depression ($\beta = 0.002$, 95 % LBCI 0.0005 to 0.0046), accounting for 14.59 % and 9.94 % of the variance of IL-6 and depressive symptoms, respectively. Due to the cross-sectional nature of assessment of inflammation and depression findings should be approached with caution; however, results suggest that complex interactions of psychoneuroimmunological and metabolic factors underlie the association between ACEs and adulthood depression.

1. Introduction

Prevalence of adverse childhood experiences (ACEs) is remarkably high. Estimates showed that 47 % to 64 % of Western adults reported at least one ACE, while 9 % to 17 % reported four or more ACEs (Bellis et al., 2014; Swedo et al., 2023). ACEs may include exposure to several types of abuse and neglect in childhood, including physical, emotional, and sexual abuse and physical and emotional neglect (Bernstein et al., 1998). Strong evidence demonstrated that exposure to ACEs is associated with higher risk of negative psychosocial/behavioural and medical outcomes (Petrucelli et al., 2019). ACEs have been particularly associated with a higher risk of adulthood depression, with ORs ranging from 1.34 to 3.17 in meta-analysis (e.g., Tan and Mao, 2023). The underlying psychobiological factors linking ACEs to depressive symptomatology, however, are still elusive, with inflammatory response being a putative mechanism (Job et al., 2020; Maayan and Maayan, 2023).

Inflammation is considered a primary defence of the body against

physical damage, pathogen exposure, and psychosocial threats (e.g., Hänsel et al., 2010) which is orchestrated by innate immune system. According to the social signal transduction theory of depression (Slavich and Irwin, 2014; Shields et al., 2024), experiences of social threat and adversity upregulate components of immune system involved in inflammation that may drive the pathogenesis of depression. Consistently, a previous meta-analysis showed that peripheral levels of proinflammatory cytokines such as interleukin-6 (IL-6) and tumour necrosis factor- α (TNF- α) and acute phase inflammatory proteins such as C-reactive protein (CRP) were higher in individuals exposed to ACEs than in non-exposed controls (Baumeister et al., 2016), although more recent evidence found predominantly non-significant associations between ACEs and elevated levels of all inflammatory markers in adulthood (Brown et al., 2021). Traditional mechanisms proposed to be involved in ACEs-related inflammation include sympathetic nervous system activation following exposure to early life stress which may determine a shift in the profile of innate immune cells with an increase

* Corresponding author at: Department of Psychology, Faculty of Medicine and Psychology, Sapienza University of Rome, Via dei Marsi 78, 00185 Rome, Italy.
E-mail address: andrea.ballesio@uniroma1.it (A. Ballesio).

<https://doi.org/10.1016/j.jad.2024.04.072>

Received 9 February 2024; Received in revised form 5 April 2024; Accepted 21 April 2024

Available online 25 April 2024

0165-0327/© 2024 The Author(s). Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

long-term upregulation of proinflammatory gene transcription in monocytes (e.g., Cole et al., 2012; Mondelli and Vernon, 2019); greater methylation of the glucocorticoid receptors (Klengel et al., 2013); increased nuclear factor-kappa B activity, a key transcriptional control pathway in the inflammatory response (Pace et al., 2012); and glucocorticoids resistance, i.e., the inability of cortisol to exert its anti-inflammatory action (Miller et al., 2002).

Notably, while normal, temporally restricted activation of inflammatory response is not harmful *per se*, chronically or disproportionately activated inflammation may become a pathogenetic factor for physical illness (Furman et al., 2019) and could be associated with the onset of depressive symptoms in a subgroup of individuals (Milaneschi et al., 2021). Indeed, strong evidence supports the presence of higher levels of CRP, IL-6, and TNF- α in patients with depression compared to healthy controls (Dowlati et al., 2010; Osimo et al., 2020), although this is not fully explained by the increased presence of ACEs in this population (Pitharouli et al., 2021). Levels of inflammatory markers may also prospectively predict the severity of depression (Mac Giollabhui et al., 2021; Ballesio et al., 2024). In healthy samples, experimental endotoxemia, a translational model of systemic inflammation has been associated with depressogenic effects (e.g., Lasselin et al., 2021). Finally, recent studies using Mendelian randomization, an analytic procedure using genetic variants which is not susceptible to confounding and reverse causality bias (Ballesio, 2023), suggested that genetically determined inflammation was associated with depression (Kappelmann et al., 2021; Galan et al., 2022).

From this perspective, it is plausible that the association between ACEs and adult depression may be, at least partially, mediated by inflammation. This hypothesis, despite being widely accepted (Slavich and Irwin, 2014; Baumeister et al., 2016; Maayan and Maayan, 2023), is still lacking meta-analytic investigation, which is considered fundamental in evidence-based healthcare (Gopalakrishnan and Ganeshkumar, 2013). Meta-analytic structural equation modelling is an advanced statistical technique which enables to test mediation hypotheses in systematic reviews, increasing statistical power and precision of estimate compared to bivariate meta-analysis, ultimately enriching the robustness of the findings (Zagaria et al., 2023). Also, meta-analytic structural equation modelling allows to account for potential confounders and test sequential mediation (i.e., with two or more mediators). This represents an advancement compared to traditional pairwise meta-analysis, where confounders are predominantly examined as moderators through meta-regressions in an attempt to explain between-study heterogeneity. On the other hand, meta-analytic structural equation modelling permits disentangling the unique contribution of each predictor after controlling for potential confounding factors (Cheung and Hong, 2017). An accurate control of confounders appears of utmost importance in this field, as the associations between ACEs, inflammation, and depression may be remarkably influenced by several factors including body mass index (BMI, e.g., Danese and Tan, 2014; Mitchell et al., 2018; Brown et al., 2021) and age (Maayan and Maayan, 2023). To advance the field, we therefore conducted a systematic review and structural equation modelling meta-analysis testing the association between ACEs and depression and the mediating role of inflammation.

2. Materials and methods

The study followed the updated Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines (Page et al., 2021) and was registered to the PROSPERO International database of the University of York Centre for Reviews (ID number CRD42023471274).

2.1. Search strategy and selection

A systematic literature search of published studies was conducted using Pubmed, Scopus, Medline, Psycinfo, and CINAHL. Additional

records were searched by consulting a previous meta-analysis on ACEs and inflammation (Baumeister et al., 2016). The search string included the following keywords: Childhood Maltreatment OR Childhood Trauma OR Childhood Adversity OR Early Life Stress OR Child Abuse OR Child Neglect OR Adverse Childhood Experiences AND C-reactive Protein OR CRP OR Tumour Necrosis Factor OR TNF- α OR Cytokine OR Interleukin OR IL-6 OR Inflammatory OR Inflammation AND depressi*. Database search was conducted by the first and the second authors under the supervision of the last author on 2 October 2023. Articles were included if they provided original data about the association (e.g., correlation coefficients, effect sizes of group differences) of any ACE before age 18 with peripheral markers of inflammation and depressive symptoms/diagnosis in adult humans (individuals aged 18 or older). Several biomarkers of inflammation were eligible, including cytokines (e.g., IL-6, IL-1 β , IL-8, TNF- α , interferon- γ), chemokines, CRP, fibrinogen, and white blood cell count. The first and the second authors independently screened titles and abstracts as well as full-text articles against eligibility criteria. Disagreements between the reviewers were resolved by discussion with the last author. Detailed search strategy for each database is reported in the Supplementary Material.

2.2. Data extraction and quality appraisal

Data from included studies were extracted by the second author using a standardized datasheet form built to extract the following study characteristics: number of participants, % of females, mean age, type of sample (i.e., healthy individuals, patients with depression), type of inflammatory marker, depression measure, ACE measure. Also, correlation coefficients between ACEs, inflammatory markers, and depression were extracted to proceed with meta-analytic calculations. If correlation coefficients were not available in original studies, they were derived from other effect size metrics (e.g., Cohen's *d*, Odds Ratio) following Lipsey and Wilson (2001) and Borenstein et al. (2009). Alternatively, *t*-value and or *F*-ratio statistics were transformed into Cohen's *d* and then converted into correlation coefficients (Lipsey and Wilson, 2001). Extracted data were cross-checked by the first and the last author.

The quality of the included studies was assessed using the Joanna Briggs Institute (JBI) Critical Appraisal Checklist for Analytical Cross-Sectional Studies (Moola et al., 2017). The checklist included eight statements (e.g., "Were the study subjects and the setting described in detail?", "Were the outcomes measured in a valid and reliable way?"), with four possible response options: yes, no, unclear, not applicable. The total number of "yes" answers is considered an indicator of higher study quality (Moola et al., 2017).

2.3. Data analysis

Data were analysed via R version 4.2.2 using the *metaSEM* (Cheung, 2015) and *OpenMx* (Boker et al., 2011) packages.

Meta-analytic structural equation modelling was applied through a two-stage structural equation modelling (TSSEM) approach (Cheung and Chan, 2005). In Stage 1 of the TSSEM, the correlation matrices extracted from primary studies were combined using either fixed or random effects models via maximum likelihood estimation. Specifically, the homogeneity of the correlation matrices was primarily examined within a fixed effects model. Data were considered homogeneous when the following goodness-of-fit indices met their respective cut-offs (Browne and Cudeck, 1993; Wang and Wang, 2019): the root mean square error of approximation (RMSEA; reasonable when <0.08), the comparative fit index and the Tucker-Lewis index (CFI and TLI, respectively; acceptable when >0.90), and the standardized root mean squared residual (SRMR; acceptable when <0.08). If the assumption of homogeneity proved untenable, random effects models were carried out and the heterogeneity of the correlation coefficients was quantified using the I^2 statistics (Higgins and Thompson, 2002). I^2 values of 0.25, 0.50 and 0.75 indicate low, medium, and high levels of heterogeneity,

respectively (Higgins et al., 2003). In Stage 2 of the TSSEM, the hypothesised path model was fitted on the pooled correlation matrix estimated from Stage 1 by minimising the weighted least squares fit function (Browne, 1984). Three distinct partial mediation models were examined, each exploring the role of a different inflammation marker as a mediator in the relationship between ACEs and depressive symptoms: CRP as a mediator (Model 1); IL-6 as a mediator (Model 2); TNF- α as a mediator (Model 3); composite inflammation as a mediator (Model 4). For significance testing of the indirect effects, 95 % likelihood-based confidence intervals were computed (95 % LBCI; Neale and Miller, 1997; Cheung, 2009). Furthermore, two control analyses were carried out examining the role of age and BMI as potential confounders in the mediation models.

Methods for assessing publication bias within the meta-analytic

structural equation modelling framework are still in the early stages of development (Scherer et al., 2019; Jak and Cheung, 2020). Nonetheless, exploratory analyses for assessing publication bias were conducted at the level of individual correlations, rather than on correlation matrices (e.g., Scherer et al., 2019). First, the correlations were transformed into Fisher's z and combined using random effects models via restricted maximum likelihood estimation (Borenstein et al., 2009). Thereafter, the evidence of publication bias for single correlations was examined by visually inspecting the funnel plot and by conducting Egger's regression test to detect its asymmetry (Egger et al., 1997). In the absence of any missing studies, the funnel plot's shape should resemble an inverted funnel, symmetrical in nature, with a wide base containing smaller studies exhibiting significant variability, and a narrow apex (Tehrani and Yamini, 2020).




Fig. 1. PRISMA 2020 flow diagram for new systematic reviews which included searches of databases and registers only.

From: Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ* 2021;372:n71. doi: 10.1136/bmj.n71.

For more information, visit: <http://www.prisma-statement.org/>.

3. Results

3.1. Search process

Detailed search flow is reported in Fig. 1. Database search identified 1565 records (Pubmed, $n = 317$, Scopus $n = 639$, Psycinfo, $n = 230$, CINAHL, $n = 68$, Medline, $n = 311$). One additional record (Lowry et al., 2022) was detected by consulting the meta-analysis of Baumeister et al. (2016). After duplicates removal, 679 titles and abstracts were screened against eligibility criteria. Thereafter, 114 full texts were screened, and 22 met the eligibility criteria and were included in the review.

3.2. Study characteristics

Detailed information of included studies are reported in Table 1. The mean age of the participants was 38.56 ± 8.1 years. The overall percentage of females was 67.99 %. Almost all the studies used standardized measures to investigate adverse childhood experiences (ACEs). Specifically, eight studies (Congio et al., 2022; Hostinar et al., 2017; Jones et al., 2023; Kleih et al., 2022; McCormack et al., 2021; Mehta et al., 2022; O'Shields et al., 2022; Rengasamy et al., 2022) used the

Childhood Trauma Questionnaire (CTQ) (Bernstein et al., 2003) and one (Davis et al., 2019) used the Childhood Trauma Questionnaire-Short Form (CTQ-SF) (Bernstein et al., 2003), three studies (Chiang et al., 2017; Runsten et al., 2014; Taylor et al., 2006) used the Risky Families questionnaire (RFQ) (Taylor et al., 2004), two studies (De Punder et al., 2018; Kuhlman et al., 2020) used the Early Trauma Inventory (ETI) (Bremner et al., 2000), one (Gardhouse et al., 2021) used the Stress and Adversity Inventory for Adults (STRAIN) (Slavich and Shields, 2018), one (John-Henderson et al., 2020) the Adverse Childhood Experiences Study Questionnaire (Felitti et al., 1998), one (Lowry et al., 2022) used the Life History Interview (LHI) and another one (Walker et al., 2022) the Childhood Maltreatment Interview Schedule-Short Form (CMIS-SF; Briere, 1992). Among the twenty-one studies, four used ad hoc questionnaires on ACEs. In particular, one study (Batsika et al., 2021) used 29 questions, one study (Hostinar et al., 2015) used a summary measure of adverse events experienced before age 18, one study (Nakamura et al., 2022) used a measure of the occurrence of three adverse events linked to health disparities and one study (Watt et al., 2020) used ten questions from the original ACE study (Felitti et al., 1998).

With respect to inflammatory markers, twelve studies examined IL-6 (Chiang et al., 2017; Davis et al., 2019; De Punder et al., 2018;

Table 1
Study characteristics.

Study name	N total	Female %	Mean age $\pm$ SD	Type of sample	Inflammatory marker	Depression measure	ACEs measure
Bitsika et al., 2021	221	58.8 %	44.06 $\pm$ 17.83	Healthy subjects	CRP	SDS	Ad hoc questionnaire
Chiang et al., 2017	91	57.14 %	18.37 $\pm$ 0.51	Healthy subjects	IL-6	CES-D	RFQ
Congio et al., 2022	67	nr	nr	Clinical (MDD)	hs-CRP	HDRS17	CTQ
Davis et al., 2019	770	55 %	53.50 $\pm$ 7.24	Healthy subjects	IL-6	MHI	CTQ-SF
De Punder et al., 2018	88	nr	nr	Clinical (MDD)	IL-6, WBC	MADRS	ETI
Gardhouse et al., 2021	64	100 %	28.7 $\pm$ 9.1	Clinical (MDD; BPD)	IL-6; TNF- α ; CRP	BDI-II	STRAIN
Hostinar et al., 2015	1180	56 %	57.3 $\pm$ 11.5	Healthy subjects	Composite inflammation (IL-6; CRP; fibrinogen; E-Selectin; ICAM-1)	CES-D	Ad hoc questionnaire
Hostinar et al., 2017	314	55.7 %	55.3 $\pm$ 11.2	Healthy subjects	Composite inflammation (CRP; IL-6; fibrinogen)	CES-D	CTQ
John-Henderson et al., 2020	90	50 %	37.55 $\pm$ 15.74	Healthy subjects	IL-6; CRP	PHQ-9	ACESQ
Jones et al., 2023	331	50.5 %	40.24 $\pm$ 6.24	Healthy subjects	IL-6; CRP	BDI-II	CTQ
Kleih et al., 2022	180	100 %	27.6 $\pm$ 5.2	Pregnant women	Composite inflammation (TNF- α ; IL-6)	CES-D	CTQ
Kuhlman et al., 2020	41	73.2 %	18.49 $\pm$ 0.75	Healthy subjects	IL-6	POMS-15	ETI
Lowry et al., 2022	3029	54.8 %	nr	Healthy subjects	hs-CRP	CES-D	LHI
Mehta et al., 2022	54	100 %	39.8 $\pm$ 11.90	Healthy subjects	IL-6, CRP	BDI-II	CTQ
McCormack et al., 2021	187	100 %	29.64 $\pm$ 6.25	Pregnant women	IL-6	HAM-D	CTQ
Nakamura et al., 2022	3416	58.28 %	68.41 $\pm$ 10.24	Healthy subjects	hs-CRP	CES-D	Ad hoc questionnaire
O'Shields et al., 2022	2118	54.91 %	53.02 $\pm$ 12.55	Healthy subjects	IL-6; TNF- α ; CRP, fibrinogen	CES-D	CTQ
Rengasamy et al., 2022	74	65 %	36.24 $\pm$ 10.83	Clinical (TRD)	IL-6; TNF- α	MADRS	CTQ
Runsten et al., 2014	116	nr	42.89 $\pm$ 8.09	Healthy subjects	hs-CRP	MINI	RFQ
Taylor et al., 2006	3248	nr	40.1 $\pm$ 3.6	Healthy subjects	CRP	CES-D	RFQ
Walker et al., 2022	200	61.5 %	20.49 $\pm$ 3.28	Healthy subjects	IL-6	CES-D	CMIS-SF
Watt et al., 2020	93	73 %	21.04 $\pm$ 2.05	Healthy subjects	CRP	PHQ-9	Ad hoc questionnaire

Abbreviations: ACESQ = Adverse Childhood Experiences Study Questionnaire, BDI-II = Beck Depression Inventory-II, BPD = borderline personality disorder, CES-D = Center for Epidemiologic Studies-Depression Scale, CMIS-SF = Childhood Maltreatment Interview Schedule-Short Form, CRP = C-reactive protein, CTQ = Childhood Trauma Questionnaire, CTQ-SF = Childhood Trauma Questionnaire-Short Form, ETI = Early Trauma Inventory, HAM-D = Hamilton Depression Rating Scale, HDRS17 = 17-item Hamilton Depression Rating Scale, hs-CRP = High-sensitivity C-reactive protein, ICAM-1 = Intercellular Adhesion Molecule-1, IL-6 = Interleukin 6, LHI = Life History Interview, MADRS = Montgomery-Åsberg Depression Rating Scale, MDD = major depressive disorder, MHI = Mental Health Inventory, MINI = Mini-International Neuropsychiatric Interview, NR = not reported, PHQ-9 = Patient Health Questionnaire-9, POMS-15 = Profile of Mood States, RFQ = Risky Families questionnaire, SDS = 20-item Zung Self-rating Depression Scale, STRAIN = Stress and Adversity Inventory for Adults, TNF- α = Tumour Necrosis Factor α , TRD = treatment-resistant depression, WBC = white blood cell count.

Gardhouse et al., 2021; John-Henderson et al., 2020; Jones et al., 2023; Kuhlman et al., 2020; McCormack et al., 2021; Mehta et al., 2022; O'Shields et al., 2022; Rengasamy et al., 2022; Walker et al., 2022), thirteen CRP (Bitsika et al., 2021; Congio et al., 2022; De Punder et al., 2018; Gardhouse et al., 2021; John-Henderson et al., 2020; Jones et al., 2023; Lowry et al., 2022; Mehta et al., 2022; Nakamura et al., 2022; O'Shields et al., 2022; Runsten et al., 2014; Taylor et al., 2006; Watt et al., 2020), three TNF- α (Gardhouse et al., 2021; O'Shields et al., 2022; Rengasamy et al., 2022), one white blood cell count (De Punder et al., 2018) and one fibrinogen (O'Shields et al., 2022). Although included in the systematic review, studies regarding fibrinogen and white blood cell count were not meta-analysed due to the limited amount of data. Furthermore, three studies employed measures of composite inflammation, consisting of IL-6, CRP, fibrinogen, E-Selectin and Intercellular Adhesion Molecule-1 (ICAM-1) (Hostinar et al., 2015), CRP, IL-6 and fibrinogen (Hostinar et al., 2017), and TNF- α and IL-6 (Kleih et al., 2022).

With respect to depression assessment, nine studies (Chiang et al., 2017; Hostinar et al., 2015; Hostinar et al., 2017; Kleih et al., 2022; Lowry et al., 2022; Nakamura et al., 2022; O'Shields et al., 2022; Taylor et al., 2006; Walker et al., 2022) used the Center for Epidemiologic Studies-Depression Scale (CES-D) (Radloff, 1977), while two (De Punder et al., 2018; Rengasamy et al., 2022) administered the Montgomery-Åsberg Depression Rating Scale (MADRS) (Davidson et al., 1986), two other studies (John-Henderson et al., 2020; Watt et al., 2020) used the Patient Health Questionnaire-9 (PHQ-9) (Spitzer et al., 2000), three (Gardhouse et al., 2021; Jones et al., 2023; Mehta et al., 2022) the Beck Depression Inventory-II (Beck et al., 1996), one study (Bitsika et al., 2021) used the 20-item Zung Self-rating Depression Scale (SDS) (Zung, 1965), another one (Congio et al., 2022) the 17-item Hamilton Depression Rating Scale (HDRS17) (Hamilton, 1960), one study (Davis et al., 2019) the Mental Health Inventory (MHI) (Veit and Ware, 1983), one (Kuhlman et al., 2020) used three items of the Profile of Mood States (POMS-15) (McNair et al., 1992), one (McCormack et al., 2021) the Hamilton Depression Rating Scale 21-item version (HAM-D) (Hamilton, 1960) and one (Runsten et al., 2014) the Mini-International Neuropsychiatric Interview (MINI) (Sheehan et al., 1998).

3.3. Study quality

According to the JBI Checklist for Analytical Cross-Sectional Studies (Moola et al., 2017), the maximum achievable score for each study was eight. Only three studies obtained the maximum score (De Punder et al., 2018; Gardhouse et al., 2021; Rengasamy et al., 2022). However, in eighteen studies, Item 4 was not applicable. Thirteen studies received a score of seven, three studies scored a six, two studies scored a five and one study scored a four. No studies received a score below four. Summarising qualitative strengths, most of the included studies showed a clear definition of inclusion criteria in the sample, along with a detailed description of study subjects and settings. Additionally, all the studies employed a valid and reliable measure for the outcome, along with the use of appropriate statistical analyses. In terms of weakness, the main critical points were the identification of potential confounding factors and the implementation of strategies to deal with these confounders.

3.4. Structural equation modelling meta-analysis

3.4.1. CRP as a mediator between ACEs and depressive symptoms

Stage 1 analysis based on a fixed effects TSSEM approach rejected the hypothesis of homogeneity of the correlation coefficients: $\chi^2 = 143.874$ (36), $p < .001$, CFI = 0.699; TLI = 0.674; RMSEA = 0.054 (90 % CI 0.045 to 0.064), SRMR = 0.048. Accordingly, heterogeneity was assumed and a random effects TSSEM was carried out. Table 2 shows the pooled correlation matrix along with heterogeneity statistics ($k = 13$, $n = 12,935$). The three weighted correlations were significant and in the expected direction, with I^2 statistics ranging between 0.397 and 0.898.

Table 2
Pooled correlations between study variables and heterogeneity statistics from the first stage of two-stage structural equation modelling.

	Estimate	SE	p-Value	I ²
Model 1. CRP as a mediator				
ACEs $\Leftrightarrow$ CRP	0.046	0.022	.039	0.661
ACEs $\Leftrightarrow$ Depressive symptoms	0.203	0.037	<.001	0.898
CRP $\Leftrightarrow$ Depressive symptoms	0.077	0.016	<.001	0.397
Model 2. IL-6 as a mediator				
ACEs $\Leftrightarrow$ IL-6	0.060	0.023	.009	0.270
ACEs $\Leftrightarrow$ Depressive symptoms	0.310	0.032	<.001	0.632
IL-6 $\Leftrightarrow$ Depressive symptoms	0.067	0.015	.001	0 ^b
Model 3. TNF- α as a mediator				
ACEs $\Leftrightarrow$ TNF- α	0.138	0.020	<.001	0 ^b
ACEs $\Leftrightarrow$ Depressive symptoms	0.188	0.020	<.001	0 ^b
TNF- α $\Leftrightarrow$ Depressive symptoms	0.082	0.109	.451	0.781
Model 4. Composite inflammation as a mediator ^a				
ACEs $\Leftrightarrow$ Composite inflammation	0.081	0.024	<.001	
ACEs $\Leftrightarrow$ Depressive symptoms	0.308	0.022	<.001	
Composite inflammation $\Leftrightarrow$ Depressive symptoms	0.143	0.023	<.001	

Abbreviations. ACEs: adverse childhood experiences, CRP: C-reactive protein, IL-6: interleukin-6, SE: standard error, TNF: tumour necrosis factor.

^a Heterogeneity statistics not applicable for fixed effects models.
^b The between-study variance was fixed to foster optimization and convergence.

In Stage 2, the pooled correlation matrix based on the random-effects approach was employed to fit the partial mediation model. Specifically, ACEs were significantly associated with CRP ($\beta = 0.046$, $p = .039$), which in turn was positively related to depressive symptoms ($\beta = 0.068$, $p < .001$). ACEs also exerted a direct effect on depressive symptoms ($\beta = 0.200$, $p < .001$). Moreover, CRP partially mediated the relationship between ACEs and depressive symptoms ($\beta = 0.003$, 95 % LBCI 0.0002 to 0.0068). The model accounted for 0.22 % and 4.6 % of the variance of CRP and depressive symptoms, respectively.

3.4.2. IL-6 as a mediator between ACEs and depressive symptoms

Stage 1 analysis based on a fixed effects TSSEM approach rejected the hypothesis of homogeneity of the correlation coefficients: $\chi^2 = 91.592$ (33), $p < .001$, CFI = 0.834; TLI = 0.819; RMSEA = 0.072 (90 % CI 0.054 to 0.089), SRMR = 0.070. Consistently, heterogeneity was assumed and a random effects TSSEM was conducted. Table 2 reports the pooled correlation matrix along with heterogeneity statistics ($k = 12$, $n = 4108$). All the correlation coefficients were statistically significant, with I^2 statistics ranging from 0.270 to 0.632.

In Stage 2 of the TSSEM, the pooled correlation matrix based on the random effects approach was used as the input matrix for the partial mediation model. Findings revealed that ACEs were significantly associated with IL-6 ($\beta = 0.060$, $p = .009$), which in turn was positively related to depressive symptoms ($\beta = 0.048$, $p = .002$). The direct effect of ACEs on depressive symptoms was statistically significant ($\beta = 0.307$, $p < .001$). Moreover, IL-6 partially mediated the relationship between ACEs and depressive symptoms ($\beta = 0.003$, 95 % LBCI 0.001 to 0.006). The model accounted for 0.36 % and 9.85 % of the variance of IL-6 and depressive symptoms, respectively.

3.4.3. TNF- α as a mediator between ACEs and depressive symptoms

In Stage 1, the fixed effects TSSEM did not support the homogeneity assumption: $\chi^2 = 23.911$ (6), $p = .005$, CFI = 0.872; TLI = 0.809; RMSEA = 0.063 (90 % CI 0.037 to 0.090), SRMR = 0.021. Table 2 reports the pooled correlation matrix ($k = 3$, $n = 2256$) along with I^2 statistics. ACE exhibited a significant correlation with both TNF- α and

depressive symptoms, but there was no significant correlation between TNF- α and depressive symptoms.

In Stage 2, the pooled correlation matrix based on the random effects approach was employed to fit the partial mediation model. Findings showed that ACEs were positively associated with TNF- α ($\beta = 0.138$, $p < .001$). In contrast, TNF- α showed no significant association with depressive symptoms ($\beta = 0.057$, $p = .606$). The direct path between ACEs and depressive symptoms was significant ($\beta = 0.180$, $p < .001$). Lastly, TNF- α did not mediate the relationship between ACEs and depressive symptoms ($\beta = 0.008$, 95 % LBCI -0.023 to 0.040). The model accounted for 1.91 % and 3.88 % of the variance of TNF- α and depressive symptoms, respectively.

3.4.4. Composite inflammation measure as a mediator between ACEs and depressive symptoms

In Stage 1, the fixed effects TSSEM supported the homogeneity assumption: $\chi^2 = 4.263$ (6), $p = .658$, CFI = 1.000; TLI = 1.013 RMSEA = 0.000 (90 % CI 0.000 to 0.045), SRMR = 0.018. Table 2 reports the pooled correlation matrix ($k = 3$, $n = 1674$), in which all the estimates were statistically significant and in the expected direction.

In Stage 2, a fixed effects TSSEM was specified on the pooled correlation matrix to estimate the path model. Specifically, ACEs were significantly associated with composite inflammation ($\beta = 0.081$, $p < .001$), which in turn was positively related to depressive symptoms ($\beta = 0.119$, $p < .001$). ACEs also exerted a direct effect on depressive symptoms ($\beta = 0.298$, $p < .001$). Furthermore, composite inflammation partially mediated the relationship between ACEs and depressive symptoms ($\beta = 0.009$, 95 % LBCI 0.004 to 0.018). The model accounted for 0.67 % and 10.89 % of the variance of composite inflammation and depressive symptoms, respectively.

3.4.5. Control analyses

Two control analyses were carried out considering age and BMI as covariates in the mediation models. Although covariates were not reported in all included studies, the TSSEM approach handles missing correlation coefficients through maximum likelihood estimation, which is unbiased and efficient under ignorable missing data conditions (see Cheung and Cheung, 2016).

3.4.5.1. Age as a covariate. Eight studies measuring CRP reported zero-order correlations between age and the investigated variables (Bitsika et al., 2021; De Punder et al., 2018; John-Henderson et al., 2020; Jones et al., 2023; Mehta et al., 2022; Nakamura et al., 2022; O'Shields et al., 2022; Runsten et al., 2014). When controlling for age, the mediation role of CRP in the relationship between ACEs and depressive symptoms remained statistically significant ($\beta = 0.003$, 95 % LBCI 0.0001 to 0.0069). Age did not exert any significant effect on CRP and depressive symptoms ($p > .05$).

Moreover, ten studies measuring IL-6 reported zero-order correlations between age and the investigated variables (Chiang et al., 2017; De Punder et al., 2018; John-Henderson et al., 2020; Jones et al., 2023; Kuhlman et al., 2020; McCormack et al., 2021; Mehta et al., 2022; O'Shields et al., 2022; Rengasamy et al., 2022; Walker et al., 2022). When controlling for age, the indirect effect of ACEs on depressive symptoms through IL-6 vanished ($\beta = 0.003$, 95 % LBCI -0.0007 to 0.0094). More specifically, age was significantly associated with IL-6 ($\beta = 0.064$, $p = .038$) but not with depressive symptoms ($p > .05$).

Unfortunately, we were unable to run sensitivity analyses on composite inflammation ($k = 2$) and TNF- α ($k = 2$) due to the limited data.

3.4.5.2. BMI as a covariate. Seven studies measuring CRP reported zero-order correlations between BMI and the investigated variables (De Punder et al., 2018; John-Henderson et al., 2020; Jones et al., 2023; Mehta et al., 2022; Nakamura et al., 2022; O'Shields et al., 2022; Taylor et al., 2006). When controlling for BMI, the indirect effect between ACEs

and depressive symptoms via CRP vanished ($\beta = 0.002$, 95 % LBCI -0.001 to 0.005). More specifically, BMI was marginally associated with ACEs ($\beta = 0.048$, $p = .053$), as well as significantly associated with CRP ($\beta = 0.350$, $p < .001$). This resulted in a non-significant relationship between ACEs and CRP ($\beta = 0.029$, $p = .208$).

Moreover, nine studies measuring IL-6 reported zero-order correlations between BMI and the investigated variables (Chiang et al., 2017; De Punder et al., 2018; John-Henderson et al., 2020; Jones et al., 2023; Kuhlman et al., 2020; McCormack et al., 2021; Mehta et al., 2022; O'Shields et al., 2022; Rengasamy et al., 2022). When controlling for BMI, IL-6 did not mediate the relationship between ACEs and depressive symptoms ($\beta = 0.001$, 95 % LBCI -0.002 to 0.004). More specifically, BMI was positively associated with ACEs ($\beta = 0.107$, $p < .001$) and IL-6 ($\beta = 0.379$, $p < .001$). This resulted in a non-significant relationship between ACEs and IL-6 ($\beta = 0.019$, $p = .429$).

Unfortunately, we were unable to run sensitivity analyses on composite inflammation ($k = 1$) and TNF- α ($k = 2$) due to the limited data.

3.4.5.3. Sequential mediation model. On the basis of such results and from a data-driven perspective, we tested a sequential mediation of BMI and IL-6 in the path between ACEs and depressive symptoms (Fig. 2). Findings revealed a significant relationship between ACEs and BMI ($\beta = 0.107$, $p < .001$), as well as between BMI and IL-6 ($\beta = 0.379$, $p < .001$) and between IL-6 and depressive symptoms ($\beta = 0.050$, $p = .009$). Moreover, ACEs exerted a direct effect on depressive symptoms ($\beta = 0.308$, $p < .001$). Most importantly, the indirect effect linking ACEs to IL-6 via BMI was significant ($\beta = 0.041$, 95 % LBCI 0.024 to 0.063), as well as the sequential mediating effect of BMI and IL-6 in the relationship between ACEs and depressive symptoms ($\beta = 0.002$, 95 % LBCI 0.0005 to 0.0046). Overall, the model accounted for 14.59 % and 9.94 % of the variance of IL-6 and depressive symptoms, respectively.

As a further step, a subgroup analysis was conducted by excluding a single sample of pregnant women (McCormack et al., 2021). Findings supported the sequential mediation of BMI and IL-6 in the relationship between ACEs and depressive symptoms ($\beta = 0.002$, 95 % LBCI 0.0002 to 0.0039). Additionally, we conducted a further subgroup analysis by considering only healthy samples. More specifically, three studies (Rengasamy et al., 2022; De Punder et al., 2018; Gardhouse et al., 2021) conducted on clinical samples with mood disorders were excluded from the analysis. Although the effect remained consistent in size, the sequential mediation linking ACEs to depressive symptoms via BMI and IL-6 was no longer significant, possibly due to lower power ($\beta = 0.002$, 95 % LBCI -0.001 to 0.004). Furthermore, we performed a leave-one-out sensitivity analysis, wherein multiple meta-analyses were conducted on each subset of the studies obtained by excluding one study at a time. Findings revealed that the sequential mediation effect did not demonstrate robustness under a null hypothesis significance testing approach (see Supplementary Document). Specifically, while the sequential mediation effect remained significant when excluding 7 out of the 12 studies included in the meta-analysis, it lost significance when excluding 5 out of the 12 studies. This may be partially attributed to lower statistical power, as the effect sizes remained consistent throughout each step of the analysis.

Finally, considering that both the assessment of the mediator and the outcome occurred relatively at the same time, we proceeded to investigate the feasibility of an alternative mediation pathway. In fact, previous research has highlighted that depressive symptoms may predict obesity and higher BMI (e.g., Franko et al., 2005), also underscoring the bidirectional relationship between inflammation and depressive episodes (e.g., Messay et al., 2012; Mac Giollabhui et al., 2021). As detailed in the Supplementary Document, the alternative sequential mediation model linking ACEs to IL-6 through depression symptoms and BMI was not supported by the data ($\beta = 0.001$, 95 % LBCI -0.004 to 0.006).




Fig. 2. Sequential mediation model. * $p < .01$, ** $p < .001$. Abbreviations: ACEs, adverse childhood experiences; BMI, body mass index; IL-6, interleukin-6.

4. Publication bias

Publication bias was assessed for each single correlation reported by at least ten studies (Higgins and Green, 2011). The visual inspection of the funnel plot indicated a reasonable degree of symmetry (see Supplementary Document). Moreover, Egger's tests were not significant for all the examined correlations: ACEs with CRP (1.829, $p = .067$), CRP with depressive symptoms (-1.571 , $p = .116$); ACE with IL-6 (0.077, $p = .938$); IL-6 with depressive symptoms (0.091, $p = .927$). The only exception was the association between ACEs and depressive symptoms, which showed a significant degree of asymmetry (3.013, $p = .003$; see Supplementary Fig. 5). Overall, these findings suggested a minimal impact of publication bias on the examined correlations.

5. Discussion

The present meta-analysis indicates that the inflammatory markers, CRP, IL-6, TNF- α and a composite inflammation measure, exert mediation effects in the relationship between ACEs and adult depressive symptoms which are explained by the strong association between BMI and inflammation. Thus, meta-analytic results provide evidence for a complex model, with ACEs predicting adult depression via the sequential mediation of BMI and IL-6. Precisely, the model showed that ACEs predict higher BMI, which in turn predicts higher IL-6, which eventually predicts depressive symptoms severity.

A slight yet significant increase in the risk of overweight and obesity for individuals exposed to ≥ 2 ACEs has been reported in previous meta-analyses of children and adult samples (Elsenburg et al., 2017; Wiss and Brewerton, 2020; Hughes et al., 2017), with several putative underlying mechanisms (Wiss and Brewerton, 2020). Experiencing physical or sexual abuse in childhood and/or adolescence can predict the development of food addiction in adulthood (Mason et al., 2016), which is a well-known determinant of obesity (Pedram et al., 2013). Moreover, it has been suggested that the high inflammatory response to stress associated with child abuse may reinforce overeating and craving for palatable foods (Hemmingsson, 2018), which might serve as self-medication/compensation mechanisms against trauma (Mason et al., 2016). Interestingly, previous research has indicated that exposure to ACEs has long-term effects on low fruit and vegetable consumption, resulting in unhealthy dietary choices, through its detrimental impact on self-regulation skills (Horino and Yang, 2021; Miller et al., 2011). In this regard, ACEs have demonstrated a significant role in predicting adults' emotional dysregulation (Rudenstine et al., 2019), which is crucial in predisposing risk for overweight and obesity (Leehr et al., 2015; Shriver et al., 2019). The current study, in combination with prior research, highlighted the need for future analyses to explore these mechanisms and their potential interactions in explaining the association between ACEs and BMI.

The mediatory role of BMI between ACEs and inflammation is consistent with previous evidence. Petrov et al. (2016) reported that BMI significantly mediated the association between childhood emotion, physical and sexual abuse and CRP, IL-6, and fibrinogen; Mitchell et al. (2018) showed that BMI mediated the relationship between physical abuse and both serum CRP and IL-6, but not TNF- α . Maes et al. (2018) found that the significant effects of early life trauma on CRP levels were completely mediated by BMI. Summarising previous evidence, however, an updated narrative review recently concluded that evidence on the mediation of BMI between childhood trauma and inflammation was substantially inconclusive (Brown et al., 2021). As a major progression from Brown et al. (2021) review, our meta-analysis, albeit limited by the cross-sectional nature of the data, suggests that the mediation of BMI between ACEs and inflammation is indeed plausible, and can be further, sequentially linked to depression. Of note, this path was detected for IL-6 but not for CRP, likely due to the smaller association between BMI and ACEs in the studies testing the CRP mediation model. Supporting our findings, the alternative pathway linking ACEs to inflammation through depression symptoms and BMI was not significant in our cross-sectional analysis.

Possible immune-to-brain pathways leading to depression have been previously extensively reviewed (e.g., Zunszain et al., 2013; Osimo et al., 2020; Nettis and Pariante, 2020; Ballesio, 2023). These include cytokine production in the brain following microglial activation through humoral (i.e., cytokines diffusion across intact or leaky regions of the blood-brain-barrier, i.e., circumventricular organs), neuronal (e.g., binding of peripheral cytokines to afferent vagus nerve fibres), or cellular pathways (Bower and Kuhlman, 2023), exacerbating changes in monoaminergic neurotransmission, neurotrophic activity, and oxidative stress processes with the following onset of cognitive, affective and motivational symptoms of depression (e.g., Bakunina et al., 2015; Dantzer, 2018; Nettis and Pariante, 2020; Turkheimer et al., 2023; Ballesio, 2023; Ballesio et al., 2023). Consistent with such hypotheses, case-control studies detected the presence of higher corticospinal fluid concentrations of inflammatory markers including IL-6 in patients with depression compared to controls (Enache et al., 2019), and experimental intravenous endotoxin injection in healthy humans induces upregulation of serum and cerebrospinal fluid concentrations of proinflammatory cytokines as well as depressive symptoms (e.g., Lasselin et al., 2021). Alternatively, it has been proposed that peripheral inflammation may reduce blood-brain-barrier permeability which in the long term may disrupt brain homeostasis and trigger depressive symptoms (Turkheimer et al., 2021).

Refining aforementioned immune-to-brain pathways, results of the present study suggest that several neuroimmunometabolic factors leading to depression could be involved in individuals with ACEs. Dietary patterns characterised by high fat intake can lead to increased amount of lipids in adipocytes (Ferreira et al., 2014), which may

increase the release of monocyte chemoattractant protein-1 (MCP-1), a chemokine that increases the infiltration of macrophages and T cells into white adipose tissue and stimulates peripheral release of pro-inflammatory cytokines including IL-6 (Shelton and Miller, 2010), finally exacerbating a state of low-grade chronic inflammation (Capuron et al., 2017). Alterations of gut microbiota, intestinal dysbiosis, and increased gut permeability may also lead to the production of peripheral inflammatory factors (Torres-Fuentes et al., 2017; Furman et al., 2019). Such immunometabolic alterations may not be restricted to periphery but rather reach the brain. The impact of diet-related inflammation in the brain has been mostly investigated at the level of hypothalamus. For instance, Lopes et al. (2024) recently showed that high fat diet for four weeks resulted in body weight gain and adiposity as well as increased gene expression of pro-inflammatory cytokines such as IL-6 and dysregulation in the nuclear factor kappa B pathway in the hypothalamus of male mice. Notably, such processes have been hypothesised to be involved in hypothalamic circuitry remodelling (Miller and Spencer, 2014), dysregulate HPA axis activity and induce subsequent production of glucocorticoids (Burfeind et al., 2016), and eventually determine depressive symptoms (Cernackova et al., 2020). Moreover, another recent mice model suggested that alternation of microbiota may be associated with microglial activation by modulating RAS-NF- κ B signal pathway (Liu et al., 2024). It was beyond the aim of the current meta-analysis to clarify the precise biological processes linking ACEs, BMI, IL-6, and depressive symptoms, that should be further elucidated in future mechanistic research.

5.1. Clinical implications

Present results may implicate that BMI and inflammation could be a potential target for depression prevention in individuals with a history of trauma or ACEs. The finding that the association between ACEs and inflammation could be best described as indirect via BMI may suggest that effective clinical interventions focused on early life exposures to adversities and subsequent inflammation should include the assessment of weight gain trajectories. Parallely, early trauma-informed screening may be successful in reducing the risk of overweight/obesity (Wiss and Brewerton, 2020), thus indirectly protecting against associated inflammation. Nevertheless, clinicians should also contemplate the evidence that BMI could be indicative of multiple latent health-related conditions (e.g., cardiovascular disease, diabetes, mood disorders) in order to appropriately address its relationships with cumulative childhood stress exposure. In this respect, the identification of pernicious behavioural mechanisms (e.g., overeating) and potential protective factors (e.g., emotion regulation) underlying the ACEs-BMI pathway should be an important public health priority as it would be informative in delineating the best strategies for the prevention and treatment of inflammation in individuals with trauma histories.

Notably, cumulative evidence has indicated the type and the severity of traumatic events as the second most associated determinant of treatment-seeking, after psychopathology itself (Gavrilovic et al., 2005). Previous research suggested that about 37 % of individuals with a history of ACEs seek psychological support (Sheerin et al., 2016), although the type of intervention offered to these patients rarely takes into consideration its impact on biological variables such as inflammation. Research highlighted significant broader deficiencies in psychological support approaches for victims of traumatic events associated with comorbid chronic inflammation (Glynn et al., 2022). This treatment gap emphasized the urge to implement integrated healthcare approaches to address medical and psychological comorbidities accompanying ACEs and the underlying mechanism of inflammation. We recently conducted a comprehensive network meta-analysis on the impact of psychotherapies on immune outcomes suggesting that mindfulness-based cognitive therapy and mindfulness-based stress reduction interventions may consistently be associated with reduced inflammation in randomised controlled trials (Ballesio et al., 2023). However, whether this reduction

may drive improvements in depression, especially in individuals with ACEs is still not demonstrated. Further research on how these biological processes, associated with psychological support of individuals with ACEs, could affect depression aetiology is imperative.

5.2. Limitations

Several limitations should be mentioned. First, the assessment of putative mediator (i.e., inflammation) and outcome (i.e., depression) occurred relatively at the same time. With cross-sectional data, we cannot infer that changes in inflammatory levels that are associated with ACEs predict changes in depressive symptoms. To thoroughly investigate a longitudinal mediation hypothesis, the prior levels of a variable must be measured and controlled for (see Little, 2013); therefore, the directionality of these associations remains to be confirmed using multiple assessments (e.g., Zagaria et al., 2023). Related to this, multiple assessment design would allow to fully test an alternative pathway linking ACEs to inflammation through depression symptoms and BMI which, as mentioned above, was not significant in our cross-sectional analysis (see Supplemental Material). Second, several variables that are presumed to play a key role were not considered in the present study due to a lack of data needed for meta-analytic calculations. Namely, ACEs are associated with physical inactivity, smoking, heavy alcohol use, and diabetes (Hughes et al., 2017), which could all be associated also with BMI. Greater childhood trauma has also been significantly associated with poorer sleep (Brindle et al., 2018), which is a strong predictor of depression (Hertenstein et al., 2019) and which may increase inflammatory signalling (e.g., Ballesio et al., 2022; Zagaria et al., 2022) and BMI (Norton et al., 2018; Lombardo et al., 2020). In order to consider these factors in further structural equation modelling meta-analyses, future studies are encouraged to report zero-order correlation coefficients with the main variables (ACEs, inflammation, depression). Third, the composite inflammation score incorporated different inflammatory biomarkers and its interpretation might be biased by the heterogeneity of indicators used for its computation (Strawbridge et al., 2015). To date, which biomarkers best represent a comprehensive index of inflammation is under debate in the literature. The identification of an optimal approach to operationalizing the composite inflammation score requires further investigation. Finally, due to limited data available, we were unable to consider the differential impact of different types of ACEs (e.g., parental absence, sexual abuse, physical abuse; Lacey et al., 2020), nor several potential other moderators such as duration of ACEs exposure, the context of the stressor, and relationship to the perpetrator (Baumeister et al., 2016).

6. Conclusion

Regardless of the aforementioned limitations, the present meta-analysis has the strength to provide at least two novel insights into the study of depressogenic mechanisms associated with ACEs: 1) it explains why the available literature on the role of inflammation as a single mediator between ACEs and depression yielded only weak and spurious effects; 2) it suggests that BMI is not a mere confounder, but rather a plausible process variable in the association between ACEs, inflammation, and depressive symptoms. Putatively, precise biological (e.g., HPA axis activity, neurotrophic factors, oxidative stress processes) and psychological (e.g., affect regulation, health behaviour, eating patterns) mechanisms are likely to play a role and should be integrated into even more complex models. Moreover, different path directions (i.e., ACEs leading to higher BMI through the mediation of depressive symptoms) are also plausible and should be investigated. Finally, we would also point out that ACEs may lead to several other psychopathological outcomes including post-traumatic stress disorders, psychotic disorders, and personality disorders (Hughes et al., 2017; Cerrato et al., 2017) in which depression is often comorbid. An effort should be made to explore potential disease-specific mechanisms linking ACEs to these conditions.

However, our study confirms the crucial role of inflammation in the paths between ACEs and depression, even as one element of complex sequential models.

Funding/acknowledgement

None.

CRediT authorship contribution statement

Andrea Zagaria: Writing – review & editing, Writing – original draft, Methodology, Formal analysis, Data curation. **Valeria Fiori:** Writing – review & editing, Writing – original draft, Methodology, Data curation. **Mariacarla Vacca:** Writing – review & editing, Writing – original draft, Methodology, Investigation. **Caterina Lombardo:** Writing – original draft. **Carmine M. Pariante:** Writing – review & editing, Writing – original draft. **Andrea Ballesio:** Writing – review & editing, Writing – original draft, Supervision, Methodology, Investigation, Conceptualization.

Declaration of generative AI and AI-assisted technologies in the writing process

Authors did not use generative AI and AI-assisted technologies in the writing this manuscript.

Declaration of competing interest

Authors have no conflict of interest to disclose.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.jad.2024.04.072>.

References

- Bakunina, N., Pariante, C.M., Zunszain, P.A., 2015. Immune mechanisms linked to depression via oxidative stress and neuroprogression. *Immunology* 144 (3), 365–373 ([doi:10.1111/imm.12443](https://doi.org/10.1111/imm.12443)).
- Ballesio, A., 2023. Where does inflammation in insomnia come from? and does it matter for comorbidity? *Sleep* 46 (10), zsad223. <https://doi.org/10.1093/sleep/zsad223>.
- Ballesio, A., Zagaria, A., Ottaviani, C., Steptoe, A., Lombardo, C., 2022. Sleep disturbance, neuro-immune markers, and depressive symptoms in older age: conditional process analysis from the English Longitudinal Study of Aging (ELSA). *Psychoneuroendocrinology* 142, 105770. <https://doi.org/10.1016/j.psyneuen.2022.105770>.
- Ballesio, A., Zagaria, A., Vacca, M., Pariante, C.M., Lombardo, C., 2023. Comparative efficacy of psychological interventions on immune biomarkers: a systematic review and network meta-analysis (NMA). *Brain Behav. Immun.* 111, 424–435. <https://doi.org/10.1016/j.bbi.2023.05.006>.
- Ballesio, A., Micheli, F., Baccini, F., Zagaria, A., Del Forno, A., Fiori, V., Lombardo, C., 2024. Inflammation as an aetiological trigger for depressive symptoms in a prospective cohort of patients with inflammatory bowel disease. *J. Psychosom. Res.* 177, 111592. <https://doi.org/10.1016/j.jpsychores.2024.111592>.
- Batsika, V., Sharpley, C.F., McMillan, M.E., Jesulola, E., Agnew, L.L., 2021. Effects of subtypes of child maltreatment on CRP in adulthood. *Front. Psychiatry* 12, 533722. <https://doi.org/10.3389/fpsy.2021.533722>.
- Baumeister, D., Akhtar, R., Ciufolini, S., Pariante, C.M., Mondelli, V., 2016. Childhood trauma and adulthood inflammation: a meta-analysis of peripheral C-reactive protein, interleukin-6 and tumour necrosis factor- α . *Mol. Psychiatry* 21 (5), 642–649. <https://doi.org/10.1038/mp.2015.67>.
- Beck, A.T., Steer, R.A., Brown, G., 1996. Beck depression inventory–II. *Psychol. Assess.* <https://doi.org/10.1037/t00742-000>.
- Bellis, M.A., Hughes, K., Leckenby, N., Perkins, C., Lowey, H., 2014. National household survey of adverse childhood experiences and their relationship with resilience to health-harming behaviors in England. *BMC Med.* 12 (1), 1–10. <https://doi.org/10.1186/1741-7015-12-72>.
- Bernstein, D.P., Fink, L., Handelsman, L., Foote, J., 1998. Childhood trauma questionnaire. In: *Assessment of Family Violence: A Handbook for Researchers and Practitioners*.
- Bernstein, D.P., Stein, J.A., Newcomb, M.D., Walker, E., Pogge, D., Ahluvalia, T., Stokes, J., Handelsman, L., Medrano, M., Desmond, D., Zule, W., 2003. Development and validation of a brief screening version of the Childhood Trauma Questionnaire. *Child Abuse Negl.* 27 (2), 169–190. [https://doi.org/10.1016/S0145-2134\(02\)00541-0](https://doi.org/10.1016/S0145-2134(02)00541-0).
- Bitsika, V., Sharpley, C.F., McMillan, M.E., Jesulola, E., Agnew, L.L., 2021. Effects of subtypes of child maltreatment on CRP in adulthood. *Front. Psych.* 12, 533722. <https://doi.org/10.3389/fpsy.2021.533722>.
- Boker, S., Neale, M., Maes, H., Wilde, M., Spiegel, M., Brick, T., Spies, J., Estabrook, R., Kenny, S., Bates, T., Mehta, P., Fox, J., 2011. OpenMx: an open source extended structural equation modeling framework. *Psychometrika* 76, 306317. <https://doi.org/10.1007/s11336-010-9200-6>.
- Borenstein, M., Hedges, L.V., Higgins, J.P., Rothstein, H.R., 2009. *Introduction to Meta-Analysis*. John Wiley & Sons, Ltd.
- Bower, J.E., Kuhlman, K.R., 2023. Psychoneuroimmunology: an introduction to immune-to-brain communication and its implications for clinical psychology. *Ann. Rev. Clin. Psychol.* 9 (19), 331–359. <https://doi.org/10.1146/annurev-clinpsy-080621-045153>.
- Bremner, J.D., Vermetten, E., Mazure, C.M., 2000. Development and preliminary psychometric properties of an instrument for the measurement of childhood trauma: the Early Trauma Inventory. *Depress. Anxiety* 12 (1), 1–12. [https://doi.org/10.1002/1520-6394\(2000\)12:1<1::AID-DA1>3.0.CO;2-W](https://doi.org/10.1002/1520-6394(2000)12:1<1::AID-DA1>3.0.CO;2-W).
- Briere, J.N., 1992. *Child Abuse Trauma: Theory and Treatment of the Lasting Effects*. Sage Publications, Inc.
- Brindle, R.C., Cribbet, M.R., Samuelsson, L.B., Gao, C., Frank, E., Krafty, R.T., Thayer, J. F., Buysse, D.J., Hall, M.H., 2018. The relationship between childhood trauma and poor sleep health in adulthood. *Psychosom. Med.* 80 (2), 200. <https://doi.org/10.1097/PSY.0000000000000542>.
- Brown, M., Worrell, C., Pariante, C.M., 2021. Inflammation and early life stress: an updated review of childhood trauma and inflammatory markers in adulthood. *Pharmacol. Biochem. Behav.* 211, 173291. <https://doi.org/10.1016/j.pbb.2021.173291>.
- Browne, M.W., 1984. Asymptotically distribution-free methods for analysis of covariance structures. *Br. J. Math. Stat. Psychol.* 37, 62–83.
- Browne, M.W., Cudeck, R., 1993. Alternative ways of assessing model fit. In: Bollen, K.A., Long, J.S. (Eds.), *Testing Structural Equation Models*. Sage Publications, Newbury Park, CA, pp. 136–162.
- Burfeind, K.G., Michaelis, K.A., Marks, D.L., 2016. The central role of hypothalamic inflammation in the acute illness response and cachexia. *Semin. Cell Dev. Biol.* 54, 42–52. <https://doi.org/10.1016/j.semcdb.2015.10.038>.
- Capuron, L., Lassel, J., Castanon, N., 2017. Role of adiposity-driven inflammation in depressive morbidity. *Neuropsychopharmacology* 42 (1), 115–128. <https://doi.org/10.1038/npp.2016.123>.
- Cernackova, A., Durackova, Z., Trebatcka, J., Mravec, B., 2020. Neuroinflammation and depressive disorder: the role of the hypothalamus. *J. Clin. Neurosci. Off. J. Neurosurgical Soc. Australasia* 75, 5–10. <https://doi.org/10.1016/j.jocn.2020.03.005>.
- Cerrato, F., Aluisio, G., Casadio, P., Di Girolamo, G., Valente, S., De Ronchi, D., Atti, A., 2017. Adverse childhood experiences, personality disorders and addiction: which relationship? *Eur. Psychiatry* 41 (S1), S200. <https://doi.org/10.1016/j.eurpsy.2017.01.2147>.
- Cheung, M.W.L., 2009. Comparison of methods for constructing confidence intervals of standardized indirect effects. *Behav. Res. Methods* 41 (2), 425–438. <https://doi.org/10.3758/BRM.41.2.425>.
- Cheung, M.W.L., 2015. metaSEM: an R package for meta-analysis using structural equation modeling. *Front. Psychol.* 5, 1521. <https://doi.org/10.3389/fpsyg.2014.01521>.
- Cheung, M.W.L., Chan, W., 2005. Meta-analytic structural equation modeling: a two-stage approach. *Psychol. Methods* 10 (1), 40–64. <https://doi.org/10.1037/1082-989X.10.1.40>.
- Cheung, M.W., Cheung, S.F., 2016. Random-effects models for meta-analytic structural equation modeling: review, issues, and illustrations. *Res. Synth. Methods* 7 (2), 140–155. <https://doi.org/10.1002/jrsm.1166>.
- Cheung, M.W.L., Hong, R.Y., 2017. Applications of meta-analytic structural equation modelling in health psychology: examples, issues, and recommendations. *Health Psychol. Rev.* 11 (3), 265–279. <https://doi.org/10.1080/17437199.2017.1343678>.
- Chiang, J.J., Bower, J.E., Irwin, M.R., Taylor, S.E., Fuligni, A.J., 2017. Adiposity moderates links from early adversity and depressive symptoms to inflammatory reactivity to acute stress during late adolescence. *Brain Behav. Immun.* 66, 146–155. <https://doi.org/10.1016/j.bbi.2017.06.015>.
- Cole, S.W., Conti, G., Arevalo, J.M., Ruggiero, A.M., Heckman, J.J., Suomi, S.J., 2012. Transcriptional modulation of the developing immune system by early life social adversity. *Proc. Natl. Acad. Sci. U. S. A.* 109 (50), 20578–20583. <https://doi.org/10.1073/pnas.1218253109>.
- Congio, A.C., Urbano, M.R., Soares, M.R.Z., Nunes, S.O.V., 2022. Cognitive impairment, childhood trauma, sedentary behaviour, and elevated C-reactive protein levels in major affective disorders. *J. Psychiatr. Res.* 155, 1–9. <https://doi.org/10.1016/j.jpsychires.2022.07.052>.
- Danese, A., Tan, M., 2014. Childhood maltreatment and obesity: systematic review and meta-analysis. *Mol. Psychiatry* 19 (5), 544–554. <https://doi.org/10.1038/mp.2013.54>.
- Dantzer, R., 2018. Neuroimmune interactions: from the brain to the immune system and vice versa. *Physiol. Rev.* 98 (1), 477–504. <https://doi.org/10.1152/physrev.00039.2016>.
- Davidson, J., Turnbull, C.D., Strickland, R., Miller, R., Graves, K., 1986. The Montgomery-Åsberg Depression Scale: reliability and validity. *Acta Psychiatr. Scand.* 73 (5), 544–548. <https://doi.org/10.1111/j.1600-0447.1986.tb02723.x>.
- Davis, M.C., Lemery-Chalfant, K., Yeung, E.W., Lueken, L.J., Zautra, A.J., Irwin, M.R., 2019. Interleukin-6 and depressive mood symptoms: mediators of the association between childhood abuse and cognitive performance in middle-aged adults. *Ann. Behav. Med.* 53 (1), 29–38. <https://doi.org/10.1093/abm/kay014>.

- De Punder, K., Entringer, S., Heim, C., Deuter, C.E., Otte, C., Wingenfeld, K., Kuehl, L.K., 2018. Inflammatory measures in depressed patients with and without a history of adverse childhood experiences. *Front. Psych.* 9, 610. <https://doi.org/10.3389/fpsy.2018.00610>.
- Dowlati, Y., Herrmann, N., Swardfager, W., Liu, H., Sham, L., Reim, E.K., Lanctôt, K.L., 2010. A meta-analysis of cytokines in major depression. *Biol. Psychiatry* 67 (5), 446–457. <https://doi.org/10.1016/j.biopsych.2009.09.033>.
- Egger, M., Smith, G.D., Schneider, M., Minder, C., 1997. Bias in meta-analysis detected by a simple, graphical test. *Bmj* 315 (7109), 629–634. <https://doi.org/10.1136/bmj.315.7109.629>.
- Elsenburg, L.K., van Wijk, K.J.E., Liefbroer, A.C., Smidt, N., 2017. Accumulation of adverse childhood events and overweight in children: a systematic review and meta-analysis. *Obesity (Silver Spring, Md.)* 25 (5), 820–832. <https://doi.org/10.1002/oby.21797>.
- Enache, D., Pariante, C.M., Mondelli, V., 2019. Markers of central inflammation in major depressive disorder: a systematic review and meta-analysis of studies examining cerebrospinal fluid, positron emission tomography and post-mortem brain tissue. *Brain Behav. Immun.* 81, 24–40. <https://doi.org/10.1016/j.bbi.2019.06.015>.
- Felitti, V.J., Anda, R.F., Nordenberg, D., Williamson, D.F., Spitz, A.M., Edwards, V., Marks, J.S., 1998. Relationship of childhood abuse and household dysfunction to many of the leading causes of death in adults: the Adverse Childhood Experiences (ACE) study. *Am. J. Prev. Med.* 14 (4), 245–258. [https://doi.org/10.1016/S0749-3797\(98\)00017-8](https://doi.org/10.1016/S0749-3797(98)00017-8).
- Ferreira, A.V., Menezes-Garcia, Z., Viana, J.B., Mário, E.G., Botion, L.M., 2014. Distinct metabolic pathways trigger adipocyte fat accumulation induced by high-carbohydrate and high-fat diets. *Nutrition (Burbank, Los Angeles County, Calif.)* 30 (10), 1138–1143. <https://doi.org/10.1016/j.nut.2014.02.017>.
- Franko, D.L., Striegel-Moore, R.H., Thompson, D., Schreiber, G.B., Daniels, S.R., 2005. Does adolescent depression predict obesity in black and white young adult women? *Psychol. Med.* 35 (10), 1505–1513. <https://doi.org/10.1017/S0033291705005386>.
- Furman, D., Campisi, J., Verdine, E., Carrera-Bastos, P., Targ, S., Franceschi, C., Ferrucci, L., Gilroy, G.W., Miller, A.H., Mantovani, A., Weyand, C.M., Barzilai, N., Goronz, J.J., Rando, T.A., Effros, R.B., Lucia, A., Kleinstreuer, N., Slavich, G.M., 2019. Chronic inflammation in the etiology of disease across the life span. *Nat. Med.* 25 (12), 1822–1832. <https://doi.org/10.1038/s41591-019-0675-0>.
- Galan, D., Perry, B.L., Warrier, V., Davidson, C.C., Stupart, O., Easton, D., Khandaker, G.M., Murray, G.K., 2022. Applying Mendelian randomization to appraise causality in relationships between smoking, depression and inflammation. *Sci. Rep.* 12 (1), 15041. <https://doi.org/10.1038/s41598-022-19214-4>.
- Gardhouse, K., Carcone, D., Ruocco, A.C., 2021. Stressed, sick, and sad: Neuroendocrine pathways between subjective lifetime stress and depression. *Brain Behav. Immun. Health* 14, 100249. <https://doi.org/10.1016/j.bbih.2021.100249>.
- Gavrilovic, J.J., Schützwohl, M., Fazel, M., Priebe, S., 2005. Who seeks treatment after a traumatic event and who does not? A review of findings on mental health service utilization. *J. Traumatic Stress Off. Publ. Int. Soc. Traumatic Stress Stud.* 18 (6), 595–605. <https://doi.org/10.1002/jts.20068>.
- Glynn, H., Apputhurai, P., Moore, G.T., Knowles, S.R., 2022. Exploring the impact of trauma and psychosocial processes on the quality of life in individuals living with inflammatory bowel disease. *J. Gastrointest. Liver Dis.* 31 (3) <https://doi.org/10.15403/jgld-4302>.
- Gopalakrishnan, S., Ganeshkumar, P., 2013. Systematic reviews and meta-analysis: understanding the best evidence in primary healthcare. *J. Family Med. Prim. Care* 2 (1), 9–14. <https://doi.org/10.4103/2249-4863.109934>.
- Hamilton, M., 1960. A rating scale for depression. *J. Neurol. Neurosurg. Psychiatry* 23 (1), 56. <https://doi.org/10.1136/jnnp.23.1.56>.
- Hänsel, A., Hong, S., Cámara, R.J., Von Känel, R., 2010. Inflammation as a psychophysiological biomarker in chronic psychosocial stress. *Neurosci. Biobehav. Rev.* 35 (1), 115–121. <https://doi.org/10.1016/j.neubiorev.2009.12.012>.
- Hemmingson, E., 2018. Early childhood obesity risk factors: socioeconomic adversity, family dysfunction, offspring distress, and junk food self-medication. *Curr. Obes. Rep.* 7 (2), 1–209. <https://doi.org/10.1007/s13679-018-0310-2>.
- Hertenstein, E., Feige, B., Gmeiner, T., Kienzler, C., Spiegelhalder, K., Johann, A., Jansson-Fröjmark, M., Palagini, L., Rücker, G., Riemann, D., Baglioni, C., 2019. Insomnia as a predictor of mental disorders: a systematic review and meta-analysis. *Sleep Med. Rev.* 43, 96–105. <https://doi.org/10.1016/j.smrv.2018.10.006>.
- Higgins, J.P.T., Green, S., 2011. *Cochrane handbook for systematic reviews of interventions*. Cochrane Collab.
- Higgins, J.P., Thompson, S.G., 2002. Quantifying heterogeneity in a meta-analysis. *Stat. Med.* 21 (11), 1539–1558. <https://doi.org/10.1002/sim.1186>.
- Higgins, J., Thompson, S.G., Deeks, J.J., Altman, D.G., 2003. Measuring inconsistency in meta-analysis. *Br. Med. J.* 327, 557–560. <https://doi.org/10.1136/bmj.327.7414.557>.
- Horino, M., Yang, W., 2021. Impact of adverse childhood experiences and fruit and vegetable intake in adulthood. *Public Health Nutr.* 24 (5), 1034–1041. <https://doi.org/10.1017/S1368980019004932>.
- Hostinar, C.E., Lachman, M.E., Mroczek, D.K., Seeman, T.E., Miller, G.E., 2015. Additive contributions of childhood adversity and recent stressors to inflammation at midlife: findings from the MIDUS study. *Dev. Psychol.* 51 (11), 1630. <https://doi.org/10.1037/dev0000049>.
- Hostinar, C.E., Davidson, R.J., Graham, E.K., Mroczek, D.K., Lachman, M.E., Seeman, T.E., van Reekum, C.M., Miller, G.E., 2017. Frontal brain asymmetry, childhood maltreatment, and low-grade inflammation at midlife. *Psychoneuroendocrinology* 75, 152–163. <https://doi.org/10.1016/j.psyneuen.2016.10.026>.
- Hughes, K., Bellis, M.A., Hardcastle, K.A., Sethi, D., Butchart, A., Mikton, C., Jones, L., Dunne, M.P., 2017. The effect of multiple adverse childhood experiences on health: a systematic review and meta-analysis. *Lancet Public Health* 2 (8), e356–e366. [https://doi.org/10.1016/S2468-2667\(17\)30118-4](https://doi.org/10.1016/S2468-2667(17)30118-4).
- Iob, E., Lacey, R., Steptoe, A., 2020. Adverse childhood experiences and depressive symptoms in later life: longitudinal mediation effects of inflammation. *Brain Behav. Immun.* 90, 97–107. <https://doi.org/10.1016/j.bbi.2020.07.045>.
- Jak, S., Cheung, M.W.L., 2020. Meta-analytic structural equation modeling with moderating effects on SEM parameters. *Psychol. Methods* 25 (4), 430. <https://doi.org/10.1037/met0000245>.
- John-Henderson, N.A., Henderson-Matthews, B., Ollinger, S.R., Racine, J., Gordon, M.R., Higgins, A.A., Horn, W.C., Reeves, S.A., Running Wolf, J.A., Grant, D., Rynda-Apple, A., 2020. Adverse childhood experiences and immune system inflammation in adults residing on the blackfeet reservation: the moderating role of sense of belonging to the community. *Ann. Behav. Med.* 54 (2), 87–93. <https://doi.org/10.1093/abm/kaz029>.
- Jones, E.J., Marsland, A.L., Gianaros, P.J., 2023. Do trait-level emotion regulation strategies moderate associations between retrospective reports of childhood trauma and prospective changes in systemic inflammation? *Stress and Health* 39 (3), 525–538.
- Kappelmann, N., Arloth, J., Georgakis, M.K., Czamara, D., Rost, N., Ligthart, S., Khandaker, G.M., Binder, E.B., 2021. Dissecting the association between inflammation, metabolic dysregulation, and specific depressive symptoms: a genetic correlation and 2-sample Mendelian randomization study. *JAMA Psychiatry* 78 (2), 161–170. <https://doi.org/10.1001/jamapsychiatry.2020.3436>.
- Kleith, T.S., Entringer, S., Scholaske, L., Kathmann, N., DePunder, K., Heim, C.M., Wadhwa, P.D., Buss, C., 2022. Exposure to childhood maltreatment and systemic inflammation across pregnancy: the moderating role of depressive symptomatology. *Brain Behav. Immun.* 101, 397–409. <https://doi.org/10.1016/j.bbi.2022.02.004>.
- Klengel, T., Mehta, D., Anacker, C., Rex-Haffner, M., Pruessner, J.C., Pariante, C.M., Pace, T.W., Mercer, K.B., Mayberg, H.S., Bradley, B., Nemeroff, C.B., Holsboer, F., Heim, C.M., Ressler, K.J., Rein, T., Binder, E.B., 2013. Allele-specific FKBP5 DNA demethylation mediates gene-childhood trauma interactions. *Nat. Neurosci.* 16 (1), 33–41. <https://doi.org/10.1038/nn.3275>.
- Kuhlman, K.R., Robles, T.F., Haydon, M.D., Dooley, L., Boyle, C.C., Bower, J.E., 2020. Early life stress sensitizes individuals to the psychological correlates of mild fluctuations in inflammation. *Dev. Psychobiol.* 62 (3), 400–408. <https://doi.org/10.1002/dev.21908>.
- Lacey, R.E., Pinto Pereira, S.M., Li, L., Danese, A., 2020. Adverse childhood experiences and adult inflammation: single adversity, cumulative risk and latent class approaches. *Brain Behav. Immun.* 87, 820–830. <https://doi.org/10.1016/j.bbi.2020.03.017>.
- Lasselin, J., Lekander, M., Benson, S., Schedlowski, M., Engler, H., 2021. Sick for science: experimental endotoxemia as a translational tool to develop and test new therapies for inflammation-associated depression. *Mol. Psychiatry* 26 (8), 3672–3683. <https://doi.org/10.1038/s41380-020-00869-2>.
- Leehr, E.J., Krohmer, K., Schag, K., Dresler, T., Zipfel, S., Giel, K.E., 2015. Emotion regulation model in binge eating disorder and obesity—a systematic review. *Neurosci. Biobehav. Rev.* 49, 125–134. <https://doi.org/10.1016/j.neubiorev.2014.12.008>.
- Lipsey, M.W., Wilson, D.B., 2001. *Practical Meta-analysis*. Sage Publications, Inc.
- Little, T.D., 2013. *Longitudinal Structural Equation Modeling*. Guilford Press.
- Liu, J., Lv, X., Ye, T., Zhao, M., Chen, Z., Zhang, Y., Yang, W., Xie, H., Zhan, L., Chen, L., Liu, W.C., Su, K.P., Sun, J., 2024. Microbiota-microglia crosstalk between Blautia producta and neuroinflammation of Parkinson's disease: a bench-to-bedside translational approach. *Brain Behav. Immun.* 9 <https://doi.org/10.1016/j.bbi.2024.01.010>. S0889-1591(24)00010-2.
- Lombardo, C., Ballesio, A., Gasparrini, G., Cerolini, S., 2020. Effects of acute and chronic sleep deprivation on eating behaviour. *Clin. Psychol.* 24 (1), 64–72. <https://doi.org/10.1111/cp.12189>.
- Lopes, P.K.F., Costa, S.O., Simino, L.A.P., Chaves, W.F., Silva, F.A., Costa, C.L., Milanski, M., Ignacio-Souza, L.M., Torsoni, A.S., Torsoni, M.A., 2024. Hypothalamic inflammation and the development of an obese phenotype induced by high-fat diet consumption is exacerbated in alpha7 nicotinic cholinergic receptor knockout mice. *Food Res. Int. (Ottawa, Ont.)* 176, 113808. <https://doi.org/10.1016/j.foodres.2023.113808>.
- Lowry, E., McInerney, A., Schmitz, N., Deschênes, S.S., 2022. Adverse childhood experiences and cognitive function in adulthood: examining the roles of depressive symptoms and inflammation in a prospective cohort study. *Soc. Psychiatry Psychiatr. Epidemiol.* 57 (12), 2367–2377. <https://doi.org/10.1007/s00127-022-02315-w>.
- Maayan, L., Maayan, M., 2023. Inflammatory mediation of the relationship between early adversity and major depressive disorder: a systematic review. *J. Psychiatr. Res.* <https://doi.org/10.1016/j.jpsychires.2023.11.025>.
- Mac Giollaibhui, N., Ng, T.H., Ellman, L.M., Alloy, L.B., 2021. The longitudinal associations of inflammatory biomarkers and depression revisited: systematic review, meta-analysis, and meta-regression. *Mol. Psychiatry* 26 (7), 3302–3314. <https://doi.org/10.1038/s41380-020-00867-4>.
- Maes, M., Congio, A., Moraes, J.B., Bonifácio, K.L., Barbosa, D.S., Vargas, H.O., Nunes, S.O.V., 2018. Early life trauma predicts affective phenomenology and the effects are partly mediated by staging coupled with lowered lipid-associated antioxidant defences. *Biomol. Concepts* 9 (1), 115–130. <https://doi.org/10.1515/bmc-2018-0010>.
- Mason, S.M., Austin, S.B., Bakalar, J.L., Boynton-Jarrett, R., Field, A.E., Gooding, H.C., Holsen, L.M., Jackson, B., Neumark-Sztainer, D., Sanchez, M., Sogg, S., Tanofsky-Kraff, M., Rich-Edwards, J.W., 2016. Child maltreatment's heavy toll: the need for trauma-informed obesity prevention. *Am. J. Prev. Med.* 50 (5), 646–649. <https://doi.org/10.1016/j.amepre.2015.11.004>.

- McCormack, C., Lauriola, V., Feng, T., Lee, S., Spann, M., Mitchell, A., Champagne, F., Monk, C., 2021. Maternal childhood adversity and inflammation during pregnancy: interactions with diet quality and depressive symptoms. *Brain Behav. Immun.* 91, 172–180. <https://doi.org/10.1016/j.bbi.2020.09.023>.
- McNair, D.M., Lorr, M., Droppleman, L.F., 1992. *EdITS manual for the profile of mood states (POMS)*. In: *Educational and Industrial Testing Service*.
- Mehta, N.D., Stevens, J.S., Li, Z., Fani, N., Gillespie, C.F., Ravi, M., Michopoulos, V., Felger, J.C., 2022. Inflammation, amygdala-ventromedial prefrontal functional connectivity and symptoms of anxiety and PTSD in African American women recruited from an inner-city hospital: preliminary results. *Brain Behav. Immun.* 105, 122–130. <https://doi.org/10.1016/j.bbi.2022.06.013>.
- Messay, B., Lim, A., Marsland, A.L., 2012. Current understanding of the bi-directional relationship of major depression with inflammation. *Biol. Mood Anxiety Disorders* 2, 4. <https://doi.org/10.1186/2045-5380-2-4>.
- Milaneschi, Y., Kappelmann, N., Ye, Z., Lamers, F., Moser, S., Jones, P.B., Burgess, S., Penninx, B.W.J.H., Khandaker, G.M., 2021. Association of inflammation with depression and anxiety: evidence for symptom-specificity and potential causality from UK Biobank and NESDA cohorts. *Mol. Psychiatry* 26 (12), 7393–7402. <https://doi.org/10.1038/s41380-021-01188-w>.
- Miller, A.A., Spencer, S.J., 2014. Obesity and neuroinflammation: a pathway to cognitive impairment. *Brain Behav. Immun.* 42, 10–21. <https://doi.org/10.1016/j.bbi.2014.04.001>.
- Miller, G.E., Cohen, S., Ritchey, A.K., 2002. Chronic psychological stress and the regulation of pro-inflammatory cytokines: a glucocorticoid-resistance model. *Health Psychol.* 21 (6), 531. <https://doi.org/10.1037/0278-6133.21.6.531>.
- Miller, G.E., Chen, E., Parker, K.J., 2011. Psychological stress in childhood and susceptibility to the chronic diseases of aging: moving toward a model of behavioral and biological mechanisms. *Psychol. Bull.* 137 (6), 959. <https://doi.org/10.1037/a0024768>.
- Mitchell, A.M., Porter, K., Christian, L.M., 2018. Examination of the role of obesity in the association between childhood trauma and inflammation during pregnancy. *Health Psychol.* 37 (2), 114–124. <https://doi.org/10.1037/hea0000559>.
- Mondelli, V., Vernon, A.C., 2019. From early adversities to immune activation in psychiatric disorders: the role of the sympathetic nervous system. *Clin. Exp. Immunol.* 197 (3), 319–328. <https://doi.org/10.1111/cei.13351>.
- Moola, S., Munn, Z., Tufanaru, C., Aromataris, E., Sears, K., Sfetcu, R., Currie, M., Lisy, K., Qureshi, R., Mattis, P., Mu, P., 2017. Chapter 7: Systematic reviews of etiology and risk. In: Aromataris, E., Munn, Z. (Eds.), *Joanna Briggs Institute Reviewer's Manual*. <https://doi.org/10.46658/JBIMES-20-08>. Adelaide, SA, Australia. Retrieved from: The Joanna Briggs Institute.
- Nakamura, J.S., Kim, E.S., Rentscher, K.E., Bower, J.E., Kuhlman, K.R., 2022. Early-life stress, depressive symptoms, and inflammation: the role of social factors. *Aging Ment. Health* 26 (4), 843–851. <https://doi.org/10.1080/13607863.2021.1876636>.
- Neale, M.C., Miller, M.B., 1997. The use of likelihood-based confidence intervals in genetic models. *Behav. Genet.* 27 (2), 113–120. <https://doi.org/10.1023/a:1025681223921>.
- Nettis, M.A., Pariante, C.M., 2020. Is there neuroinflammation in depression? Understanding the link between the brain and the peripheral immune system in depression. *Int. Rev. Neurobiol.* 152, 23–40. <https://doi.org/10.1016/bi.im.2019.12.004>.
- Norton, M.C., Eleuteri, S., Carolini, S., Ballesio, A., Conte, S.C., Falaschi, P., Lucidi, F., 2018. Is poor sleep associated with obesity in older adults? A narrative review of the literature. *Eat. Weight Disord.* 23 (1), 23–38. <https://doi.org/10.1007/s40519-017-0453-2>.
- O'Shields, J., Patel, D., Mowbray, O.P., 2022. Childhood maltreatment and inflammation: leveraging structural equation modeling to test the social signal transduction theory of depression. *J. Affect. Disord.* 311, 173–180. <https://doi.org/10.1016/j.jad.2022.05.077>.
- Osimo, E.F., Pillinger, T., Rodriguez, I.M., Khandaker, G.M., Pariante, C.M., Howes, O.D., 2020. Inflammatory markers in depression: a meta-analysis of mean differences and variability in 5,166 patients and 5,083 controls. *Brain Behav. Immun.* 87, 901–909. <https://doi.org/10.1016/j.bbi.2020.02.010>.
- Pace, T.W., Wingenfeld, K., Schmidt, I., Meinlschmidt, G., Hellhammer, D.H., Heim, C.M., 2012. Increased peripheral NF- κ B pathway activity in women with childhood abuse-related posttraumatic stress disorder. *Brain Behav. Immun.* 26 (1), 13–17. <https://doi.org/10.1016/j.bbi.2011.07.232>.
- Page, M.J., McKenzie, J.E., Bossuyt, P.M., Boutron, I., Hoffmann, T.C., Mulrow, C.D., Shamseer, L., Tetzlaff, J.M., Akl, E.A., Brennan, S.E., Chou, R., Glanville, J., Grimshaw, J.M., Hrobjartsson, A., Lalu, M.M., Li, T., Loder, E.W., Mayo-Wilson, E., McDonald, S., McGuinness, L.A., Moher, D., 2021. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *Int. J. Surg.* 88, 105906. <https://doi.org/10.1016/j.ijsu.2021.105906>.
- Pedram, P., Wadden, D., Amini, P., Gulliver, W., Randell, E., Cahill, F., Vasdev, S., Goodridge, A., Carter, J.C., Zhai, G., Ji, Y., Sun, G., 2013. Food addiction: its prevalence and significant association with obesity in the general population. *PLoS One* 8 (9), e7483. <https://doi.org/10.1371/journal.pone.0074832>.
- Petrov, M.E., Davis, M.C., Belyea, M.J., Zautra, A.J., 2016. Linking childhood abuse and hypertension: sleep disturbance and inflammation as mediators. *J. Behav. Med.* 39 (4), 716–726. <https://doi.org/10.1007/s10865-016-9742-x>.
- Petrucelli, K., Davis, J., Berman, T., 2019. Adverse childhood experiences and associated health outcomes: a systematic review and meta-analysis. *Child Abuse Negl.* 97, 104127. <https://doi.org/10.1016/j.chiabu.2019.104127>.
- Pitharoulis, M.C., Hagenaars, S.P., Glanville, K.P., Coleman, J.R.I., Hotopf, M., Lewis, C.M., Pariante, C.M., 2021. Elevated C-reactive protein in patients with depression, independent of genetic, health, and psychosocial factors: results from the UK biobank. *Am. J. Psychiatry* 178 (6), 522–529. <https://doi.org/10.1176/appi.ajp.2020.20060947>.
- Radloff, L.S., 1977. The CES-D scale: a self-report depression scale for research in the general population. *Appl. Psychol. Measur.* 1 (3), 385–401. <https://doi.org/10.1177/014662167700100306>.
- Rengasamy, M., Brundin, L., Griffio, A., Panny, B., Capan, C., Forton, C., Price, R.B., 2022. Cytokine and reward circuitry relationships in treatment-resistant depression. *Biol. Psychiatry Glob. Open Sci.* 2 (1), 45–53. <https://doi.org/10.1016/j.bpsgos.2021.06.009>.
- Rudenstine, S., Espinosa, A., McGee, A.B., Routhier, E., 2019. Adverse childhood events, adult distress, and the role of emotion regulation. *Traumatology* 25 (2), 124. <https://doi.org/10.1037/trm0000176>.
- Runsten, S., Korkeila, K., Koskenvuo, M., Rautava, P., Vainio, O., Korkeila, J., 2014. Can social support alleviate inflammation associated with childhood adversities? *Nord. J. Psychiatry* 68 (2), 137–144. <https://doi.org/10.3109/08039488.2013.786133>.
- Scherer, R., Siddiq, F., Tondeur, J., 2019. The technology acceptance model (TAM): a meta-analytic structural equation modeling approach to explaining teachers' adoption of digital technology in education. *Comput. Educ.* 128, 13–35. <https://doi.org/10.1016/j.compedu.2018.09.009>.
- Sheehan, D.V., Lecrubier, Y., Sheehan, K.H., Amorim, P., Janavs, J., Weiller, E., Hergueta, T., Baker, R., Dunbar, G.C., 1998. The mini-international neuropsychiatric interview (MINI): the development and validation of a structured diagnostic psychiatric interview for DSM-IV and ICD-10. *J. Clin. Psychiatry* 59 (20), 22–33.
- Sheerin, C., Berenz, E.C., Knudsen, G.P., Reichborn-Kjennerud, T., Kendler, K.S., Aggen, S.H., Amstadter, A.B., 2016. A population-based study of help seeking and self-medication among trauma-exposed individuals. *Psychol. Addict. Behav.* 30 (7), 771. <https://doi.org/10.1037/adb0000185>.
- Shelton, R.C., Miller, A.H., 2010. Eating ourselves to death (and despair): the contribution of adiposity and inflammation to depression. *Prog. Neurobiol.* 91 (4), 275–299. <https://doi.org/10.1016/j.pneurobio.2010.04.004>.
- Shields, G.S., Vinograd, M., Bui, T., Sichko, S., Irwin, M.R., Slavich, G.M., 2024. Heightened neural activity and functional connectivity responses to social rejection in female adolescents at risk for depression: Testing the Social Signal Transduction Theory of Depression. *J. Affect. Disord.* 15 (345), 467–476. <https://doi.org/10.1016/j.jad.2023.10.113>.
- Shriver, L.H., Doll, J.M., Lawless, M., Calkins, S.D., Keane, S.P., Shanahan, L., Wideman, L., 2019. Longitudinal associations between emotion regulation and adiposity in late adolescence: indirect effects through eating behaviors. *Nutrients* 11 (3), 517. <https://doi.org/10.3390/nu11030517>.
- Slavich, G.M., Irwin, M.R., 2014. From stress to inflammation and major depressive disorder: a social signal transduction theory of depression. *Psychol. Bull.* 140 (3), 774. <https://doi.org/10.1037/a0035302>.
- Slavich, G.M., Shields, G.S., 2018. Assessing lifetime stress exposure using the Stress and Adversity Inventory for Adults (Adult STRAIN): an overview and initial validation. *Psychosom. Med.* 80 (1), 17. <https://doi.org/10.1097/PSY.0000000000000534>.
- Spitzer, R.L., Williams, J.B., Kroenke, K., Hornyk, R., McMurray, J., Patient Health Questionnaire Obstetrics-Gynecology Study Group, 2000. Validity and utility of the PRIME-MD patient health questionnaire in assessment of 3000 obstetric-gynecologic patients: the PRIME-MD patient health questionnaire obstetrics-gynecology study. *Am. J. Obstet. Gynecol.* 183 (3), 759–769. <https://doi.org/10.1067/mob.2000.106580>.
- Strawbridge, R., Arnone, D., Danese, A., Papadopoulos, A., Vives, A.H., Cleare, A.J., 2015. Inflammation and clinical response to treatment in depression: a meta-analysis. *Eur. Neuropsychopharmacol.* 25 (10), 1532–1543.
- Swedo, E.A., Aslam, M.V., Dahlberg, L.L., Niolon, P.H., Guinn, A.S., Simon, T.R., Mercy, J.A., 2023. Prevalence of adverse childhood experiences among U.S. adults - behavioral risk factor surveillance system, 2011–2020. *MMWR Morb. Mortal. Wkly Rep.* 72 (26), 707–715. <https://doi.org/10.15585/mmwr.mm7226a2>.
- Tan, M., Mao, P., 2023. Type and dose-response effect of adverse childhood experiences in predicting depression: a systematic review and meta-analysis. *Child Abuse Negl.* 139, 106091. <https://doi.org/10.1016/j.chiabu.2023.106091>.
- Taylor, S.E., Lerner, J.S., Sage, R.M., Lehman, B.J., Seeman, T.E., 2004. Early environment, emotions, responses to stress, and health. *J. Pers.* 72 (6), 1365–1394. <https://doi.org/10.1111/j.1467-6494.2004.00300.x>.
- Taylor, S.E., Lehman, B.J., Kiefe, C.I., Seeman, T.E., 2006. Relationship of early life stress and psychological functioning to adult C-reactive protein in the coronary artery risk development in young adults study. *Biol. Psychiatry* 60 (8), 819–824. <https://doi.org/10.1016/j.biopsych.2006.03.016>.
- Tehrani, H.D., Yamini, S., 2020. Dataset of parenting practices, self-control and anti-social behaviors: Meta-analytic structural equation modeling. *Data Brief* 32, 106114. <https://doi.org/10.1016/j.dib.2020.106114>.
- Torres-Fuentes, C., Schellekens, H., Dinan, T.G., Cryan, J.F., 2017. The microbiota-gut-brain axis in obesity. *Lancet Gastroenterol. Hepatol.* 2 (10), 747–756. [https://doi.org/10.1016/S2468-1253\(17\)30147-4](https://doi.org/10.1016/S2468-1253(17)30147-4).
- Turkheimer, F.E., Althubaiti, N., Schubert, J., Nettis, M.A., Cousins, O., Dima, D., Mondelli, V., Bullmore, E.T., Pariante, C., Veronese, M., 2021. Increased serum peripheral C-reactive protein is associated with reduced brain barriers permeability of TSPO radioligands in healthy volunteers and depressed patients: implications for inflammation and depression. *Brain Behav. Immun.* 91, 487–497. <https://doi.org/10.1016/j.bbi.2020.10.025>.
- Turkheimer, F.E., Veronese, M., Mondelli, V., Cash, D., Pariante, C.M., 2023. Sickness behaviour and depression: an updated model of peripheral-central immunity interactions. *Brain Behav. Immun.* 111, 202–210. <https://doi.org/10.1016/j.bbi.2023.03.031>.
- Veit, C.T., Ware, J.E., 1983. The structure of psychological distress and well-being in general populations. *J. Consult. Clin. Psychol.* 51 (5), 730.

- Walker, J.H., Venta, A., Bechelli, J., Brewer, T.M., Boisvert, D., Bick, J., Lewis, R., Wells, J., Armstrong, T., 2022. Testing the role of inflammation in the relation of childhood maltreatment and suicidal ideation among young adults. *J. Clin. Psychol.* 78 (12), 2484–2496. <https://doi.org/10.1002/jclp.23347>.
- Wang, J., Wang, X., 2019. *Structural Equation Modeling: Applications Using Mplus*. John Wiley & Sons.
- Watt, T., Ceballos, N., Kim, S., Pan, X., Sharma, S., 2020. The unique nature of depression and anxiety among college students with adverse childhood experiences. *J. Child Adolesc. Trauma* 13, 163–172. <https://doi.org/10.1007/s40653-019-00270-4>.
- Wiss, D.A., Brewerton, T.D., 2020. Adverse childhood experiences and adult obesity: a systematic review of plausible mechanisms and meta-analysis of cross-sectional studies. *Physiol. Behav.* 223, 112964 <https://doi.org/10.1016/j.physbeh.2020.112964>.
- Zagaria, A., Lombardo, C., Balleio, A., 2022. Longitudinal association between sleep disturbance and inflammation, and the role of positive affect. *J. Sleep Res.* 31 (5), e13560 <https://doi.org/10.1111/jsr.13560>.
- Zagaria, A., Ottaviani, C., Lombardo, C., Balleio, A., 2023. Perseverative cognition as a mediator between perceived stress and sleep disturbance: a structural equation modeling meta-analysis (meta-SEM). *Ann. Behav. Med. Publ. Soc. Behav. Med.* 57 (6), 463–471. <https://doi.org/10.1093/abm/kaac064>.
- Zung, W.W., 1965. A self-rating depression scale. *Arch. Gen. Psychiatry* 12 (1), 63–70.
- Zunszain, P.A., Hepgul, N., Pariante, C.M., 2013. Inflammation and depression. *Curr. Top. Behav. Neurosci.* 14, 135–151. https://doi.org/10.1007/7854_2012_211.